

Literatur

1. Gielen S, Schuler G, Adams V. Cardiovascular effects of exercise training: molecular mechanisms. *Circulation* 2010;122:1221-1238
2. Hamer M. The relative influence of fitness and fatness on inflammatory factors. *Prev Med* 2007;44:3-11
3. Franson EI, Batty GD, Tabák et al. Association between changes in body composition and change in inflammatory markers: an 11-year follow-up in the Whitehall II Study. *J Clin Endocrinol Metab* 2010;95(12):5370-5374
4. Hamer M, Sabia S, Batty GD et al. Physical activity and inflammatory markers over 10 years. Follow up in men and women. From the Whitehall II Cohort Study. *Circulation* 2012;126:928-933
5. Steene-Johannessen J, Kolle E, Andersen LB et al. Adiposity, aerobic fitness, muscle fitness and markers of inflammation in children. *Med Sci Sports Exerc* 2013;Vol 45, No 4:714-721
6. Sonnenschein K, Horvath T, Müller M et al. Exercise training improves in vivo endothelial repair capacity of early endothelial progenitor cells in subjects with metabolic syndrome. *Eur J Cardiovasc Prev Rehabil* 2011;18(3):406-414
7. Andersen K, Lind L, Ingelsson E et al. Skeletal muscle morphology on risk of cardiovascular disease in elderly men. *Europ J Prev Cardiol* 2015;22(2):231-239
8. Stuart CA, McCurry MP, Marino A et al. Slow-twitch fiber proportion in skeletal muscle correlates with insulin responsiveness. *J Clin Endocrinol Metab* 2013;98:2027-2036
9. Hambrecht R, Fiehn E, Yu J et al. Effects of endurance training on mitochondrial ultrastructure and fibre type distribution in skeletal muscle of patients with stable chronic heart failure. *J Am Coll Cardiol* 1997;29:1067-1073
10. Ellison GE, Waring ChD, Vininanca et al. Pathophysiological cardiac remodeling in response to endurance exercise training: cellular and molecular mechanisms. *Heart* 2012;98:5-10
11. Nadal GB, Madhavi VM. Molecular basis of cardiac performance: Plasticity of the myocardium generated through protein isoform switches. *J Clin Invest* 1989;84:1693-1700
12. Bergmann O, Bhardwaj RD, Bernard S et al. Evidence of cardiomyocyte renewal in humans. *Science* 2009;324:98-102
13. Perkonen MA, Franco F, Lane LD et al. Cardiac atrophy after bed rest and spaceflight. *J Appl Physiol* 2001;91:645-653
14. Dorfman TA, Levine BD, Tillery T et al. Cardiac atrophy in women following bed rest. *J Appl Physiol* 2007;103:8-16
15. Rengo G, Parisi V, Femminella GD et al. Molecular aspects of the cardioprotective effects of exercise in the elderly. *Aging Clin Exp Res* 2013;25(5):487-497
16. Navarro-Arevalo A, Canavate C, Sanchez-del-Pino M. Myocardial and skeletal muscle aging and changes in oxidative stress in relationship to vigorous exercise training. *Mech Ageing Dev* 1999;108:207-217
17. Rinaldi B, Corbi G, Boccuti S et al. Exercise training affects age-induced changes in SOD and heat shock protein expression in rat heart. *Exp Gerontol* 2006;41:764-770
18. Lee Y, Min K, Talbert EE et al. Exercise protects cardiac mitochondria against ischemia-reperfusion injury. *Med Sci Sports Exerc* 2012;44(3):397-405
19. Gross GJ, Peart JN. KATP channels and myocardial preconditioning: an update. *Am J Physiol* 2003;285:H921-H930
20. French JP, Quindry JC, Falk DJ et al. Ischemia-reperfusion-induced calpain activation and SERCA2a degradation are attenuated by exercise training and calpain inhibition. *Am J Physiol* 2006;290:H128-H136
21. Fink BN, Kelly DP. Peroxisome proliferator-activated receptor γ coactivator-1 (PGC-1) regulatory cascade in cardiac physiology and disease. *Circulation* 2007;115:2540-2548

22. Lehman JJ, Barger PN, Kovacs A et al. Peroxisome proliferator-activated receptor gamma coactivator-1 promotes cardiac mitochondrial biogenesis. *J Clin Invest* 2000;106:847-856
23. Wang YX, Zhang ChL, Yu RT et al. Regulation of muscle fiber type and running endurance by PPAR δ . *PLoS Biology* 2004;2(10):1532-1539
24. Tiraby C, Langin D. PGC-1 α un co-activateur transcriptionnel impliqué dans le métabolisme. *Medicine/Science* 2005;21:49-54
25. Rasmussen P, Brassard P, Adser H et al. Evidence for a release of brain-derived neurotrophic factor from brain during exercise. *Exp Physiol* 2009;94:1062-1069
26. Erickson K, Voss MW, Prakash RS et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci USA* 2011;108:3017-3022
27. Rowe GC, Safdar A, Arany Z et al. Running forward, new frontiers in endurance exercise biology. *Circulation* 2014;129:798-810
28. Choi KM, Kim JH, Cho GJ et al. Effect of exercise training on plasma visfatin and eotaxin levels. *Europ J Endocrinol* 2007;157:437-442
29. Huang GJ, Ben-David E, Tort Piella A et al. Neurogenomic evidence for a shared mechanism of the antidepressant effects of exercise and chronic fluoxetine in mice. *PLoS One* 2012;7:e 35901
30. Hunsberger JG, Newton SS, Bennett AH et al. Antidepressant actions of the exercise-regulated gene VGF. *Nat Med* 2007;13:1476-1482
31. Moon HY, Kim SH, Yang YR et al. Macrophage migration inhibitory factor mediates the antidepressant actions of voluntary exercise. *Proc Natl Acad Sci USA* 2012;109:13094-13099
32. Shayage H, Al-Mallah MH, Hung RK et al. High exercise capacity attenuates the risk of early mortality after a first myocardial infarction: The Henry Ford Exercise Testing (FIT) Project. *Mayo Clin Proc* 2016;91(2):129-139
33. Goraya TY, Jacobsen SJ, Pellinka PA et al. Prognostic value of treadmill exercise testing in elderly persons. *Ann Intern Med* 2000;132:862-870
34. Chakravarty EF, Hubert HB, Lingula VB et al. Reduced disability and mortality among aging runners: a 21-year longitudinal study. *Arch Intern Med* 2008;168:1638-1646
35. López-Liuch G, Irusta PM, Navas P et al. Mitochondrial biogenesis and healthy aging. *Exp Gerontol* 2008;43(9):813-819
36. Safdar A, Bourgeois JM, Ogborn DI et al. Endurance exercise rescues progeroid aging and induces systemic mitochondrial rejuvenation in mtDNA mutator mice. *Proc Natl Acad Sci USA* 2011;108(10):4135-4140
37. Holloszy JO, Booth FW. Biochemical adaptations to endurance exercise in muscle. *Annu Rev Physiol* 1976;38:273-291
38. Boveris A, Navarro A. Systemic and mitochondrial adaptive responses to moderate exercise in rodents. *Free Radiac Biol Med* 2008;44:224-229
39. Hawley JA, Holloszy JO. Exercise: It's the real thing! *Nutr Rev* 2009;67:172-178
40. Rolland Y, Czerwinski S, Abellan Van Kan G et al. Sarcopenia: Its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging* 2008;12:433-450
41. Goldspink G, Fernandes KF, Williams PE et al. Age related changes in collagen gene expression in the muscles of mdx dystrophic and normal mice. *Neuromuscul Disord* 1994;4:193-191
42. Wenz T, Rossi SG, Rotundo RL et al. Increased muscle PGC-1 α expression protects from sarcopenia and metabolic disease during aging. *Proc Natl Acad Sci USA (PNAS)* 2009;106(48):20405-20410
43. Koliada AK, Krasnenkov DS, Vaiserman AM. Telomeric aging: mitotic clock or stress indicator? *Frontiers in Genetics* 2015;6:1-4
44. Song Z, Figura G, Liu Y et al. Lifestyle impacts on the aging associated expression of biomarkers of DNA damage and telomere dysfunction in human blood. *Aging Cell* 2010;9(4):607-615

45. Østhus IB, Sgura A, Berardinelli F et al. Telomere length and longterm endurance exercise: Does exercise training affect biological age. A pilot Study. *PloS One* 2012;7(12):e52769
46. Denham J, O'Brien BJ, Charchar FJ. Telomere length maintenance and cardio-metabolic prevention through exercise training. *Sports Med* 2016 Febr 25 [Epub ahead of print], PMID 26914269 [Pub Med]
47. Werner Ch, Fürster T, Widmann T et al. Physical exercise prevents cellular senescence in circulating leucocytes and in the vessel wall. *Circulation* 2009;120:2438-2447
48. Chilton WL, Marques FZ, Wert J et al. Acute exercise leads to regulation of telomere-associated genes and microRNA expression in immune cells. *PLoS One* 2014;9(4):e92088
49. Puterman E, Lin J, Blackburn E et al. The power of exercise: Buffering the effect of chronic stress on telomere length. *PLoS One* 2010;5(5):e10837
50. Morlá M, Busquets X, Pons J et al. Telomere shortening in smokers with and without COPD. *Europ Respir J* 2006;27:525-528
51. Krauss J, Farzaneh-Far R, Puterman E et al. Physical fitness and telomere length in patients with coronary heart disease: findings from the heart and soul study. *PLoS One* 2011;6(11):e26983
52. Farzaneh-Far R, Cawthon RM, Na B et al. Prognostic value of leucocyte telomere length in patients with stable coronary artery disease, data from the heart and soul study. *Arterioscler Thromb Vasc Biol* 2008;28:1379-1384
53. Brouillette S, Singh RK, Thompson JR et al. White cell telomere length and risk of premature myocardial infarction. *Arterioscler Thromb Vasc Biol* 2003;23:842-846
54. Shammass MA. Telomeres, lifestyle, cancer, and aging. *Curr Opin Clin Nutr Care* 2011;14(1):28-34
55. Willeit P, Willeit J, Mayr A et al. Telomere length and risk of incident cancer and cancer mortality. *JAMA* 2010;304(1):69-75
56. Wu X, Amos Ch, Zhu Y et al. Telomere dysfunction: a potential cancer predisposition factor. *J Natl Cancer Inst* 2003;95:1211-1218
57. Mc Grath M, Wong JYY, Michaud D et al. Telomere length, cigarette smoking, and bladder cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev* 2007;16(4):815-819
58. Neilson HK, Conroy SM, Friedenreich ChM. The influence of energetic factors on biomarkers of postmenopausal breast cancer risk. *Curr Nutr Rep* 2014;3:22-34
59. Lynch BM, Neilson HK, Friedenreich ChM. Physical activity and breast cancer prevention. *Recent Results Cancer Res* 2011;186:13-42
60. Wu Y, Zhang D, Kang S. Physical activity and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat* 2013;137(3):869-882
61. Carey AL, Steinberg GR, Macaulay SL et al. Interleukin-6 increases insulin-stimulated glucose disposal in humans and glucose uptake and fatty acid oxidation in vitro via AMP-activated protein kinase. *Diabetes* 2006;55:2688-2697
62. Wallenius V, Wallenius K, Ahren B et al. Interleukin-6-deficient mice develop mature-onset obesity. *Nat Med* 2002;8(1):75-79
63. Abbas T, Faivre E, Hölscher Ch. Impairment of synaptic plasticity and memory formation in GLP-1 receptor KO mice: Interaction between type 2 diabetes and Alzheimer's disease. *Behavioural Brain Research* 2009;205(1):265-271
64. Isacson R, Nielsen E, Dannaeus K et al. The glucagon-like peptide 1 receptor agonist exendin-4 improves reference memory performance and decreases immobility in the forced swim test. *Europ J Pharmacol* 2011;650(1):249-255
65. Boström P, Wu J, Jedrychowski MP et al. A PGC-1 α dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 2012;481:463-468
66. Timmons JA, Baar K, Davidsen PK et al. Is Irisin a human exercise gene? *Nature* 2012;488: E9-E10

67. Kim KH, Jeong YT, Oh H et al. Autophagy deficiency leads to protection from obesity and insulin resistance by inducing FGF 21 as a mitokine. *Nature Medicine* 2013;19:83-92
68. Izumiya Y, Bina HA, Ouchi N et al. FGF 21 is an Akt-regulated myokine. *FEBS Letters* 2008;582:3805-3810
69. Seldin MM, Peterson JM, Byerly MS et al. Myonectin(CTRP15), a novel myokine that links Skeletal muscle to systemic lipid homeostasis. *J Biol Chem* 2012;287(15):11968-11980
70. Gumá A, Martínez-Redondo V, López-Soldano I et al. Emergin role of neuregulin as a modulator of muscle metabolism. *Am J Physiol Endocrinol Metab* 2010;298:E742-E750
71. Broholm Ch, Laye MJ, Brandt C et al. LIF is a contraction-induced myokine stimulating human myocyte proliferation. *J Appl Physiol* 2011;111:251-259
72. Morissette MR, Stricker JC, Rosenberg MA et al. Effects of myostatin deletion in aging mice. *Aging Cell* 2009;8:573-58
73. Heinecke J, Auger-Messier M, Xu J et al. Genetic deletion of myostatin from the heart prevents skeletal muscle atrophy in heart failure. *Circulation* 2010;121(3):419-425
74. Van Berendoncks AM, Garnier A, Beckers P et al. Exercise training reverses adiponectin resistance in skeletal muscle of patients with chronic heart failure. *Heart* 2011;97:1403-1409
75. Van Berendoncks AM, Beckers P, Hoymans VY et al. Exercise training reduces circulating adiponectin levels in patients with chronic heart failure. *Clinical Science* 2010;118:281-289
76. Sofi F, Capalbo A, Cesari et al. Physical activity during leisure time and primary prevention of coronary heart disease: an updated meta-analysis of cohort studies. *Europ J Cardiovasc Prev Rehabil* 2008;15:247-257
77. Laukkanen JA, Rauramaa R, Kuri S. Exercise workload, coronary risk evaluation and the risk of cardiovascular and all-cause death in middle-aged men. *Europ J Cardiovasc Prev Rehabil* 2008;15:285-292
78. Nocon M, Hiemann T, Müller-Riemenschneider F et al. Association of physical activity with all-cause and cardiovascular mortality: a systemic review and meta-analysis. *Europ J Cardiovasc Prev Rehabil* 2008;15:239-246
79. Kodama S, Saito K, Tanaka S et al. Cardiac respiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: A meta-analysis. *JAMA* 2009;301(19):2024-2035
80. Rzehak P, Meisinger C, Woelke G et al. Weight change, weight cycling and mortality in the ERFORT Male Cohort Study. *Eur J Epidemiol* 2007;22:665-673
81. Diaz VA, Mainous AG III, Everett CJ. The association between weight fluctuation and mortality: results from a population-based cohort study. *J Commun Health* 2005;30:153-165
82. Lee D, Sui X, Artero EG et al. Longterm effects of changes in cardiorespiratory fitness and body mass index on all-cause and cardiovascular disease mortality in men. The aerobics center longitudinal study. *Circulation* 2011;124:2483-2490
83. Sorensen TI, Rissanen A, Korkeila M et al. Intention to lose weight, weight changes, and 18-y mortality in overweight individuals without co-morbidities. *PLoS Med* 2005;2:e171
84. Höggström G, Nordström A, Nordström P. High aerobic fitness in late adolescence is associated with a reduced risk of myocardial infarction later in life: a nation wide cohort study in men. *Eur Heart J* 2014(35):3133-3140
85. Berry JD, Willis B, Gupta S et al. Lifetime risks for cardiovascular disease mortality by cardiorespiratory fitness levels measured at ages 45, 55, and 65 years in men. The Cooper Center Longitudinal Study. *J Am Coll Cardiol* 2011;57:1604-1610
86. Boden WE, Franklin B, Berra K et al. Exercise as a therapeutic intervention in patients with stable ischemic heart disease: an underfilled prescription. *Am J Med* 2014;127:905-911
87. Oerkild B, Frederiksen M, Fischer Hansen J et al. Self-reported physical inactivity predicts survival after hospitalisation for heart disease. *Eur J Cardiovasc Prev Rehab* 2011;18(3):475-480

88. Martin BJ, Arena R, Haykowsky M et al. Cardiovascular fitness and mortality after contemporary cardiac rehabilitation. *Mayo Clin Proc* 2013;88(5):455-463
89. Clark AM, Hartling L, Vandermeer B et al. Meta-analysis: secondary prevention programmes for patients with coronary artery disease. *Ann Intern Med* 2005;143(9):659-672
90. Hedbäck B, Perk J, Wodlin P. Longterm reduction of cardiac mortality after myocardial infarction: 10-year results of a comprehensive rehabilitation programme. *Eur Heart J* 1993;14(6):831-835
91. Myers J, Prakash M, Froelicher V et al. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346:793-801
92. Taylor RS, Brown A, Ebrahim S et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004;116(19):682-692
93. Lee JY, Yun SC, Ahn JM et al. Impact of cardiac rehabilitation on angiographic outcomes after drug-eluting stents in patients with de novo long coronary artery lesions. *Am J Cardiol* 2014;113(12):1977-1985
94. Smith JL, Verrill TA, Boura JA et al. Effect of cardiorespiratory fitness on short-term morbidity and mortality after coronary bypass grafting. *Am J Cardiol* 2013;112(8):1104-1109
95. Hedbäck B, Perk J, Hörnblad M et al. Cardiac rehabilitation after coronary artery bypass surgery: 10-year results on mortality, morbidity and readmissions to hospital. *J Cardiovasc Risk* 2001;8(3):153-158
96. Iellamo F, Legramante JM, Massaro et al. Effects of a residential exercise training on baroreflex sensitivity and heart rate variability in patients with coronary artery disease. *Circulation* 2000;102:2588-2592
97. Seronde MF, Geha R, Puymirat E et al. Discharge heart rate and mortality after acute myocardial infarction. *Am J Med* 2014;127:954-962
98. Cornelissen VA, Onkelinx S, Goetschalckx K et al. Exercise-based cardiac rehabilitation improves endothelial function assessed by flow-mediated dilation but not by pulse amplitude tonometry. *Eur J Prev Cardiol* 2014;21(1):39-48
99. Heaps CL, Parker JL. Effects of exercise training on coronary collateralization and control of collateral resistance. *J Appl Physiol* 2011;111:587-598
100. Keating FK, Schneider DJ, Bunn JY et al. Effect of exercise training and weight loss on platelet reactivity in overweight patients with coronary artery disease. *J Cardiopulm Rehabil Prev* 2013;33(6):371-377
101. Madsen E, Videm V, Moholdt et al. Predictors of beneficial coronary plaque changes following aerobic exercise. *Med Sci Sports Exerc* 2015. Mar 13 [Epub ahead of print]
102. Kahn S, Chuturgoon AA, Naidoo DP. Telomeres and atherosclerosis. *Cardiovasc J Africa* 2012;23(10):563-571
103. Codd V, Nelson CP, Albrecht E et al. Identification of seven loci affecting mean telomere length and their association with disease. *Nat Genet* 2013;45(4):422-427
104. Werner C, Hanhoun M, Widmann T et al. Effects of physical exercise on telomere-regulating proteins, survival pathways, and apoptosis. *J Am Coll Cardiol* 2008;52:470-482
105. Cherkas LF, Hünkin JL, Kato BS et al. The association between physical activity in leisure time and leucocyte telomere length. *Arch Intern Med* 2008;168(2):154-158
106. La Rocca TJ, Seals DR, Pierce GL. Leucocyte telomere length is preserved with aging in endurance exercise-trained adults and related to maximal aerobic capacity. *Mech Ageing Dev* 2010;131(2):165-168
107. Denham J, Nelson CP, O'Brien BJ et al. Longer leucocyte telomere are associated with ultra endurance exercise independent of cardiovascular risk factors. *PLoS One* 2013; 8(7): e69377

108. Brouillette SW, Whittaker A, Stevens SE et al. Telomere length is shorter in healthy offspring of subjects with coronary artery disease: support for the telomere hypothesis. *Heart* 2008;94(4):422-425
109. Zee RYL, Michand SE, Germer S. Association of shorter mean telomere length with risk of incident myocardial infarction: a prospective, nested case-control approach. *Clinica Chimica Acta* 2009;403(1-2):139-141
110. Mainous AG III, Codd V, Diaz VA et al. Leucocyte telomere length and coronary calcification. *Atherosclerosis* 2010;210:262-267
111. Calvert PA, Liew T, Gorenne I et al. Leucocyte telomere length is associated with high-risk plaques on virtual intravascular ultrasound and increased proinflammatory activity. *Arterioscler Thromb Vasc Biol* 2011;31:2157-2164
112. Corrao G, Ghirardi A, Ibrahim B et al. Burden of new hospitalization for heart failure: a population-based investigation from Italy. *Eur J Heart Failure* 2014;16:729-736
113. Ziaeeian B, Sharma PP, Yu et al. Factors associated with variations in hospital expenditures for acute heart failure in the United States. *Am Heart J* 2015;169:282-289
114. Heidenreich PA, Albert NM, Allen LA et al. Forecasting the impact of heart failure in the United States. A policy statement from the American Heart Association. *Circ Heart Fail* 2013;6:606-619
115. Van Deursen VM, Urso R, Laroche C et al. Co-morbidities in patients with heart failure: an analysis of the European Heart Failure Pilot Survey. *Eur J Heart Fail* 2014;16:103-111
116. Swank AM, Horton J, Fleg JL et al. Modest increase in peak VO₂ is related to better clinical outcome in chronic heart failure patients. *Circ Heart Fail* 2012;5:579-585
117. Von Haehling S. The muscle in dire straits. Mechanism of wasting in heart failure. *Circulation* 2012;125:2686-2688
118. Middlekauff HR. Making the case for skeletal myopathy as the major limitation of exercise capacity in heart failure. *Circ Heart Fail* 2010;3:537-546
119. Stratton JR, Dunn JF, Adamopoulos S et al. Training partially reverses skeletal muscle metabolic abnormalities during exercise in heart failure. *J Appl Physiol* 1994;76:1575-1582
120. Adamopoulos S, Coats Aj, Brunotte F et al. Physical training improves skeletal muscle metabolism in patients with chronic heart failure. *J Am Coll Cardiol* 1993;21:1101-1106
121. Libera LD, Sabbadini R, Renken C et al. Apoptosis in the skeletal muscle of rats with HF is associated with increased serum levels of TNF- α and sphingosine. *J Mol Cell Cardiol* 2001;33:1871-1878
122. Reid MB, Li YP. Cytokines and oxidative signaling in skeletal muscle. *Acta Physiol Scand* 2001;171:225-232
123. Larsen AI, Lindal S, Aukrust P et al. Effect of exercise training on skeletal muscle fibre characteristics in men with chronic heart failure: Correlation between skeletal muscle alterations, cytokines and exercise capacity. *Int J Cardiol* 2002;83:25-32
124. Conraads VM, Beckers P, Bosmans J et al. Combined endurance/resistance training reduces plasma TNF- α receptor levels in patients with chronic heart failure and coronary artery disease. *Europ Heart J* 2002;23:1854-1860
125. Adamopoulos S, Parissis J, Karatzas et al. Physical training modulates proinflammatory cytokines and the soluble Fas/soluble Fas ligand system in patients with chronic heart failure. *J Am Coll Cardiol* 2002;39:653-663
126. Adamopoulos S, Parissis J, Kroupis et al. Physical training reduces peripheral markers of inflammation in patients with chronic heart failure. *Europ Heart J* 2001;22:791-797
127. Lenk K, Schnur R, Linke A et al. Impact of exercise training on myostatin expression in the myocardium and skeletal muscle in a chronic heart failure model. *Eur J Heart Failure* 2009;11:342-348

128. Gielen S, Adams V, Möbius-Winkler S et al. Antiinflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. *J Am Coll Cardiol* 2003;42:861-868
129. Gielen S, Adams V, Linke A et al. Exercise training in chronic heart failure: correlation between reduced local inflammation and improved oxidative capacity in the skeletal muscle. *Eur J Cardiovasc Prev Rehabil* 2008;15:43-48
130. Movahed H, Cao LB, Pitzalis M et al. Beneficial effects of exercise training in patients with chronic heart failure. *J Cardiol Therap* 2013;1:20-33
131. Giannuzzi P, Temporelli PL, Corrà U et al. Antiremodeling effects of long-term exercise training in patients with stable chronic heart failure (ELVD-CHF) trial. *Circulation* 2003;108:554-559
132. Lecker SH, Zavin A, Cao P et al. Expression of the irisin precursor FNDC5 in skeletal muscle correlates with aerobic exercise performance in patients with heart failure. *Circ Heart Fail* 2012;5:812-818
133. Francis GS, Cohn JN, Johnson G et al. Plasma norepinephrine, plasma renin activity and congestive heart failure: relation to survival and the effects of therapy in V-Heft II. *Circulation* 1993;87(Suppl 6):40-48
134. Böhm M, Swedberg K, Komajda M et al. Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *Lancet* 2010;376:886-894
135. Bonaduce D, Petretta M, Marciano F et al. Independent and incremental prognostic value of heart rate variability in patients with chronic heart failure. *Am Heart J* 1999;138:273-284
136. Magri D, Corrà U, Di Lenarda A et al. Cardiovascular mortality and chronotropic incompetence in systolic heart failure: the importance of a reappraisal of current cut-off criteria. *Eur J Heart Fail* 2014;16:201-209
137. Dunlay SM, Manemann SM, Chamberlain AM et al. Activities of daily living and outcomes in heart failure. *Circ Heart Fail* 2015;8:261-267
138. Wegrzynowska-Teodorczyk K, Rudzinska E, Lazarczyk M et al. Distance covered during a six-minute walk test predicts long-term cardiovascular mortality and hospitalisation rates in men with systolic heart failure: an observational study. *J Physiotherapy* 2013;59:177-187
139. O'Connor CM, Whellan DJ, Lee KL et al. HF-ACTION Investigators. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;301:1439-1450
140. Keteyian SJ, Leifer ES, Houston-Miller N et al. For the HF-ACTION Investigators. Relation between volume of exercise and clinical outcomes in patients with heart failure. *J Am Coll Cardiol* 2012;60:1899-1905
141. Davies EJ, Moxham T, Rees K et al. Exercise training for systolic heart failure: COCHRANE systematic review and meta-analysis. *Eur J Heart Fail* 2010;12:706-715
142. Piepoli MF, Davos C, Franis DP et al. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004;328:189-195
143. Belardinelli R, Georgiou D, Cianci G et al. Randomized controlled trial of long-term moderate exercise training in chronic heart failure. *Circulation* 1999;99:1173-1182
144. Belardinelli R, Georgiou D, Cianci G et al. 10-year exercise training in chronic heart failure. A randomized controlled trial. *J Am Coll Cardiol* 2012;60:1521-1528
145. Van der Harst P, Van der Steege G, De Boer RA et al. Telomere length of circulating leucocytes is decreased in patients with chronic heart failure. *J Am Coll Cardiol* 2007;49:1459-1464
146. Van der Harst P, Wong LS, De Boer RA et al. Possible association between telomere length and renal dysfunction in patients with chronic heart failure. *Am J Cardiol* 2008;102:207-210

147. Collerton J, Martin-Ruiz C, Kenny A et al. Telomere length is associated with left ventricular function in the oldest old: the Newcastle 857 study. *Eur Heart J* 2007;28:172-176
148. Wong LS, Huzen J, Van der Harst P et al. Anemia is associated with shorter leucocyte telomere length in patients with chronic heart failure. *Eur J Heart Failure* 2010;12:348-353
149. Yancy CW, Jessup M, Bazkurt B et al. ACCF/AHA Guideline for the management of heart failure: executive summary: a report of the ACCF/AHA Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;62(16):e147-239
150. Andersen K, Mariosa D, Adami HO et al. Dose-response relationship of total and leisure time physical activity to risk of heart failure. *Circ Heart Failure* 2014;7:701-708
151. Farrell SW, Finley CE, Radford NB et al. Cardiorespiratory fitness, body mass index, and heart failure mortality in men. *Circ Heart Fail* 2013;6:898-905
152. Khan H, Kunutsor S, Rauramaa R et al. Cardiorespiratory fitness and risk of heart failure: a population-based follow-up study. *Eur J Heart Fail* 2014;16:180-188
153. Berry JD, Pandey A, Gao A et al. Physical fitness and risk for heart failure and coronary artery disease. *Circ Heart Fail* 2013;6:627-634
154. Rahman I, Bellavia A, Wolk A et al. Relation between physical activity and heart failure risk in women. *Circ Heart Fail* 2014;7:877-881
155. Young DR, Reynolds K, Sidell M et al. Effects of physical activity and sedentary time on the risk of heart failure. *Circ Heart Fail* 2014;7:21-27
156. Ng M, Fleming T, Robinson M et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384(9945):766-781
157. McKinsey Global Institute. *Overcoming Obesity. An Initial Economic Analysis*. New York, NY: McKinsey and Company; 2014
158. Ladabaum U, Mannalithara A, Myer PA et al. Obesity, abdominal obesity, physical activity, and caloric intake in US adults: 1988 to 2010. *Am J Med* 2014;127:717-727
159. Must A, Spadano J, Coakley EH et al. The disease burden associated with overweight and obesity. *JAMA* 1999;282(16):1523-1529
160. O'Neill S, Driscoll O'. Metabolic syndrome: a closer look at the growing epidemic and its associated pathologies. *Obesity Reviews* 2015;16:1-12
161. Scuteri A, Laurant S, Cucca F et al. Metabolic syndrome across Europe: different clusters of risk factors. *Eur J Prev Cardiol* 2015;22(4):486-491
162. Clinical Guidelines on the Identification, evaluation, and treatment of overweight and obesity in adults: The evidence report: National Institutes of Health. *Obes Res* 1998; Suppl 2:51S-209S
163. Kurth T, Gaziano JM, Berger K et al. Body mass index and the risk of stroke in men. *Arch Intern Med* 2002;162:2557-2562
164. Kencanaiah S, Evans JC, Levy D et al. Obesity and the risk of heart failure. *N Engl J Med* 2002;347:305-313
165. Kannel WB, Plehn JF, Cupples LA. Cardiac failure and sudden death in the Framingham Study. *Am Heart J* 1988;115:869-875
166. El-Gamal A, Gallagher D, Nawras A et al. Effect of obesity on QT, RR, and QTc intervals. *Am J Cardiol* 1995;75:956-959
167. Poirier P, Giles TD, Bray GA et al. Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss. *Circulation* 2006;113:898-918
168. Wang YC, McPherson K, Marsh T et al. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 2011;378(9793):815-825
169. Park A, Kim Won Kon, Kwang-Hee BAE. Distinction of white, beige and brown adipocytes derived from mesenchymal stem cells. *World J Stem Cells* 2014;6(1):33-42
170. Rega-Kaun G, Kaun C, Wojta J. More than a simple storage organ: Adipose tissue as a source of adipokines involved in cardiovascular disease. *Thromb Haemost* 2013;110:641-650

171. Morelli M, Gaggini M, Daniele G et al. Ectopic fat: the true culprit linking obesity and cardiovascular disease? *Thromb Haemost* 2013;110:651-660
172. Northcott JM. Adipokines and the cardiovascular system: Mechanism mediating health and disease. *Can J Physiol Pharmacol* 2012;90:1029-1059
173. Azizi M. The effect of 8-weeks aerobic exercise training on serum leptin in untrained females. *Proc Social Behav Sciences* 2011;15:1630-1634
174. Akbarpour M. The effect of aerobic training on serum adiponectin and leptin levels and inflammatory markers of coronary heart disease in obese men. *Biol Sport* 2013;30:21-27
175. Abd El-Kader SM. Moderate versus high intensity exercise training on leptin and selected immune system response in obese subjects. *Eur J Gen Med* 2011;8(4):268-272
176. Yamauchi T, Kamon J, Minokoshi Y et al. Adiponectin stimulates glucose utilisation and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 2002;8:1288-1295
177. Matsunami T, Sato Y, Ariga S et al. Regulation of synthesis and oxidation of fatty acids by adiponectin receptors (AdipoR1/R2) and insulin receptor substrate isoforms (IRS-1/-2) of the liver in a nonalcoholic steatohepatitis animal model. *Metabolism* 2011;60:805-814
178. Okamoto Y, Arita Y, Nishida M et al. An adipocyte-derived plasma-protein, adiponectin, adheres to injured vascular walls. *Horm Metab Res* 2000;32:47-50
179. Ouchi N, Kihara S, Arita Y et al. Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF-kappa B signaling through a cAMP-dependent pathway. *Circulation* 2000;102:1296-1301
180. Kumada M, Kihara S, Ouchi N et al. Adiponectin specifically increased tissue inhibitor of metalloproteinase-1 through interleukin-10 expression in human macrophages. *Circulation* 2004;109:2046-2049
181. Ohashi K, Parker JL, Ouchi N et al. Adiponectin promotes macrophage polarization toward an antiinflammatory phenotype. *J Biol Chem* 2010;285:6153-6160
182. Ouchi N, Kihara S, Arita Y et al. Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages. *Circulation* 2001;103:1057-1063
183. Pischon T, Girman CJ, Hotamisligil GS et al. Plasma adiponectin levels and risk of myocardial infarction in men. *J Am Med Assoc* 2004;291:1730-1737
184. Persson J, Lindberg K, Gustafsson TP et al. Low plasma adiponectin concentration is associated with myocardial infarction in young individuals. *J Intern Med* 2010;268:194-205
185. Kelly KR, Navaneethan SD, Solomon TPJ et al. Lifestyle-induced decrease in fat mass improves adiponectin secretion in obese adults. *Med Sci Sports Exerc* 2014;46(5):920-926
186. Wang X, You T, Murphy K et al. Addition of exercise increases plasma adiponectin and release from adipose tissue. *Med Sci Sports Exerc* 2015; March 25 (epub ahead of print)
187. Nikseresht M, Sadeghifard N, Agha-Alinejad H et al. Inflammatory markers and adipocytokine responses to exercise training and detraining in men who are obese. *J Strength Cond Res* 2014;28(12):3399-3410
188. Arciero PJ, Baur D, Connely S et al. Timed daily ingestion of whey protein and exercise training reduces visceral adipose tissue mass and improves insulin resistance: The PRISE-Study. *J Appl Physiol* 2014;117(1):1-10
189. Jung DH, Kim YH, Kim JK et al. Relative contribution of obesity and serum adiponectin to the development of hypertension. *Diab Res Clin Pract* 2014;103:51-56
190. Yang RZ, Lee MJ, Hu H et al. Identification of omentin as a novel depot-specific adipokine in human adipose tissue: possible role in modulating insulin action. *Am J Physiol Endocrinol Metab* 2006;290:E1253-E1261
191. De Souza Batista CM, Yang RZ, Lee MJ et al. Omentin plasma levels and gene expression are decreased in obesity. *Diabetes* 2007;56:1655-1661
192. Liu R, Wang X, Bu P. Omentin-1 is associated with carotid atherosclerosis in patients with metabolic syndrome. *Diabetes Res Clin Pract* 2011;93:21-25

193. Zhong X, Li X, Liu F et al. Omentin inhibits TNF- α -induced expression of adhesion molecules in endothelial cells via ERK/NF- κ B pathway. *Biochem Biophys Res Commun* 2012;425: 401-406
194. Yamawaki H, Kuramoto J, Kameshima S et al. Omentin, a novel adipocytokine inhibits TNF α -induced vascular inflammation in human endothelial cells. *Biochem Biophys Res Commun* 2011;408:339-343
195. Shang FJ, Wang JP, Liu XT et al. Serum omentin-1 levels are inversely associated with the presence and severity of coronary artery disease in patients with metabolic syndrome. *Biomarkers* 2011;16:657-662
196. Shibata R, Ouchi N, Kikuchi R et al. Circulating omentin is associated with coronary artery disease in men. *Atherosclerosis* 2011;219:811-814
197. Wilms B, Ernst B, Gerig R et al. Plasma omentin-1 levels are related to exercise-performance in obese women and increase upon aerobic endurance training. *Exp Clin Endocrinol Diabetes* 2015;123(3):187-192
198. Saremi A, Asghari M, Ghorbani A. Effects of aerobic training on serum omentin-1 and cardiometabolic risk factors in overweight and obese men. *J Sports Sci* 2010;28(9):993-998
199. Scatena M. Osteopontin: a multifunctional molecule regulating chronic inflammation and vascular disease. *Arterioscler Thromb Vasc Biol* 2003;23:1029-1034
200. Gadeau AP, Campan M, Millet D et al. Osteopontin overexpression is associated with arterial smooth muscle cell proliferation in vitro. *Arterioscler Thromb* 1993;13:120-125
201. Lorenzen JM, Schauerte C, Hübner A et al. Osteopontin is indispensable for AP1-mediated angiotensin II-related miR-21 transcription during cardiac fibrosis. *Eur Heart J* 2015;36:2184-2196
202. Mazzali M, Kipari T, Ophascharoensuk V et al. Osteopontin—a molecule for all seasons. *QJM* 2002;95:3-13
203. Minoretto P, Falcone C, Calcagnino M et al. Prognostic significance of plasma osteopontin levels in patients with chronic stable angina. *Eur Heart J* 2006;27:802-807
204. Kato R, Momiyama Y, Ohmori R et al. High plasma levels of osteopontin in patients with restenosis after percutaneous coronary intervention. *Arterioscler Thromb Vasc Biol* 2006;26:e1-e2
205. Gomez-Ambrosi J, Catalán V, Ramirez B et al. Plasma osteopontin levels and expression in adipose tissue are increased in obesity. *J Clin Endocrinol Metab* 2007;92(9):3719-3727
206. Patel L, Buckels C, Kinghorn IJ et al. Resistin is expressed in human macrophages and directly regulated by PPAR γ activators. *Biochem Biophys Res Commun* 2003;300:472-476
207. Lehrke M, Reilly MP, Millington SC et al. An inflammatory cascade leading to hyper-resistinemia in humans. *PLoS Med* 2004;1:e45
208. Bokarewa M, Nagaev I, Dahlberg L et al. Resistin, an adipokine with potent proinflammatory properties. *J Immunol* 2005;174:5789-5795
209. Hsu WY, Chao YW, Tsai YL et al. Resistin induces monocyte-endothelial cell adhesion by increasing ICAM-1 and VCAM-1 expression in endothelial cells via p38MAPK-dependent pathway. *J Cell Physiol* 2011;226(8):2181-2188
210. Xu W, Yu L, Zhou W et al. Resistin increases lipid accumulation and CD36 expression in human macrophages. *Biochem Biophys Res Commun* 2006;351(2):376-382
211. Jamaluddin S, Weakley SM, Qizhi YQ et al. Resistin: functional roles and therapeutic considerations for cardiovascular disease. *Br J Pharmacol* 2012;165:622-632
212. Cabrera de Leon A, González DA, González AH et al. Relationships between serum resistin and fat intake, serum lipid concentrations and adiposity in the general population. *J Atheroscler Thromb* 2015;21:454-462

213. Kawamura R, Doi Y, Osawa H et al. Circulating resistin is increased with decreasing renal function in a general Japanese population: the HISAYAMA Study. *Nephrol Dial Transplant* 2010;25:3236-3240
214. Makni E, Moalla W, Benezzeddine-Boussaidi L et al. Correlation of resistin with inflammatory and cardiometabolic markers in obese adolescents with and without metabolic syndrome. *Obes Facts* 2013;6:393-404
215. Joksic J, Sopic M, Spasojevic V et al. Circulating resistin protein and mRNA concentrations and clinical severity of coronary artery disease. *BIOCHEMIA MEDICA* 2015;25(2):242-251
216. Wang H, Chen DY, Cao J et al. High serum resistin level may be an indicator of the severity of coronary disease in acute coronary syndrome. *Chin Med Sci J* 2009;24(3):161-169
217. Zhang MH, Na B, Schiller NB et al. Resistin, exercise capacity, and inducible ischemia in patients with stable coronary heart disease: data from the Heart and Soul Study. *Atherosclerosis* 2010;213(2):604-610
218. Aghapour A, Farzanegi P. Effect of six-week aerobic exercise on chemerin and resistin concentration in hypertensive postmenopausal women. *Electronic Physician* 2013;5(1):623-630
219. Jones TE, Basilio JL, Brophy PM et al. Long-term exercise training in overweight adolescents improves plasma peptide YY and resistin. *Obesity* 2009;17:1189-1195
220. Izadpanah A, Barnard RJ, Almeda AJE et al. Short-term diet and exercise intervention ameliorates inflammation and markers of metabolic health in overweight/obese children. *Am J Physiol Endocrinol Metab* 2012;303:E542-E550
221. Faber DR, de Groot PG, Visseren FL et al. Role of adipose tissue in haemostasis, coagulations and fibrinolysis. *Obes Rev* 2009;10(5):554-563
222. Hittel DS, Kraus WE, Hoffman EP. Skeletal muscle dictates the fibrinolytic state after exercise training in overweight men with characteristics of metabolic syndrome. *J Physiol* 2003;548(2):401-410
223. Keating FK, Schneider DJ, Savage PD et al. Effect of exercise training and weight loss on platelet reactivity in overweight patients with coronary artery disease. *J Cardiopulm Rehabil Prev* 2013;33(6):371-377
224. Tjønnå AE, Stølen TO, Bye A et al. Aerobic interval training reduces cardiovascular risk factors more than multitreatment approach in overweight adolescents. *Clinical Science* 2009;116:317-326
225. Ho SS, Dhaliwal SS, Hills AP et al. The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC Public Health* 2012;12:704-717
226. Schwingshackl L, Dias S, Strasser B et al. Impact of different training modalities on anthropometric and metabolic characteristics in overweight/obese subjects: a systematic review and network meta-analysis. *PLoS One* 2013;8(12):e 82853
227. Sigal RJ, Alberga AS, Goldfield GS et al. Effects of aerobic training, resistance training, or both on percentage body fat and cardiometabolic risk markers in obese adolescents: the healthy eating aerobic and resistance training in youth randomized clinical trial. *JAMA Pediatr* 2014;168:1006-1014
228. Stevens J, Cai J, Evenson KR et al. Fitness and fatness as predictors of mortality from all causes and from cardiovascular disease in men and women in the Lipid Research Clinics Study. *Am J Epidemiol* 2002;156(9):832-841229.
229. Sui X, La Monte MJ, Laditka JN et al. Cardiorespiratory fitness and adiposity as mortality predictors in older adults. *JAMA* 2007; 298(21):2507-2516
230. Ekelund U, Ward HA, Norat T et al. Physical activity and all-cause mortality across levels of overall and abdominal adiposity in European men and women: the European Prospective Investigation into Cancer and Nutrition Study (EPIC). *Am J Clin Nutr* 2015;14 Jan, publ. ahead of print doi:10.3945/ajcn.114.100065

231. Ogden CL, Carroll MD, Kit BK et al. Prevalence of obesity and trends in body mass index among US children and adolescents 1999-2010. *JAMA* 2012;307:483-490
232. Birch LL, Anzman SL. Learning to eat in an obesogenic environment: a developmental systems perspective on childhood obesity. *Child Dev Persp* 2010;4:138-143
233. Messiah SE, Vidot DC, Gumurkar S et al. Obesity is significantly associated with cardiovascular disease risk factors in 2-to 9-year-olds. *J Clin Hypertens* 2014;16:889-894
234. Wanahita N, Messerli FH, Bangalore S et al. Atrial fibrillation and obesity—results of a meta-analysis. *Am Heart J* 2008;155(2):310-315
235. Frost L, Hune LJ, Vestergaard P et al. Overweight and obesity as risk factors for atrial fibrillation or flutter: The Danish Diet, Cancer, and Health Study. *Am J Med* 2005;118:489-495
236. Abed HS, Samuel CS, Lau DH et al. Obesity results in progressive atrial structural and electrical remodeling: implications for atrial fibrillation. *Heart Rhythm* 2013;10:90-100
237. Wong CX, Abed HS, Molaei P et al. Pericardial fat is associated with atrial fibrillation severity and ablation outcome. *J Am Coll Cardiol* 2011;57(17):1745-1751
238. Mohanty S, Mohanty P, DiBiase L et al. Impact of metabolic syndrome on procedural outcomes in patients with atrial fibrillation undergoing catheter ablation. *J Am Coll Cardiol* 2012;59:1295-1301
239. Tang R, Dong J, Liu X et al. Metabolic syndrome and risk of recurrence of atrial fibrillation after catheter ablation. *Circ J* 2009;73:438-443
240. Abed HS, Nelson AJ, Richardson JD et al. Impact of weight reduction on pericardial adipose tissue and cardiac structure in patients with atrial fibrillation. *Am Heart J* 2015;169:655-662
241. Abed HS, Wittert GA, Leong DP et al. Effect of weight reduction and cardiometabolic risk factor management on syndrome burden and severity in patients with atrial fibrillation: a randomized clinical trial. *JAMA* 2013;310(19):2050-2060
242. Pathak RK, Middeldorp ME, Meredith M et al. Long-term effect of goal-directed weight management in an atrial fibrillation cohort: a long-term follow-up study (LEGACY). *J Am Coll Cardiol* 2015;65(20):2159-2169
243. Aydin AA, Kayah A, Poyraz AK et al. The relationship between coronary artery disease and pericoronary epicardial adipose tissue thickness. *J Int Med Res* 2015;43:17-25
244. Satoh M, Ishikawa Y, Takahashi Y et al. Association between oxidative DNA damage and telomere shortening in circulating endothelial progenitor cells obtained from metabolic syndrome patients with coronary artery disease. *Atherosclerosis* 2008;198(2):347-353
245. Guariguata L, Whiting DR, Hambleton I et al. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Clin Pract* 2014;103:137-149
246. Bener A, Al-Hamaq AO. Predictions burden of diabetes and economics cost: contributing risk factors of changing disease prevalence and its pandemic impact to Qatar. *Exp Clin Endocrinol Diabetes* 2016 March 16 [Epub ahead of print]
247. Zhang P, Zhang X, Brown J et al. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87:293-301
248. Tamayo T, Rosenbauer J, Wild SH et al. Diabetes in Europe: an update. *Diabetes Res Clin Pract* 2014;103:206-217
249. IDF Diabetes Atlas 6th ed. Brussels, Belgium: International Diabetes Federation; 2013
250. Beagley J, Guariguata L, Weil C et al. Global estimates of undiagnosed diabetes in adults. *Diabetes Res Clin Pract* 2014;103:150-160
251. TODAY Study Group. Rapid rise in hypertension and nephropathy in youth with type 2 diabetes: the TODAY clinical trial. *Diabetes Care* 2013;36(6):1753-1741
252. Adlise RB, D'Agostini Jr RB, Hamman RF et al. Search for Diabetes in Youth Study Group. The burden of diabetes mellitus among US youth: prevalence estimates from the Search for Diabetes in Youth Study. *Pediatrics* 2006;118(4):1510-1518

253. Patterson C, Guariguata L, Dahlquist G et al. Diabetes in the young—a global view and worldwide estimates of numbers of children with type 1 diabetes. *Diabetes Res Clin Pract* 2014;103:161-175
254. Charron MJ, Brosius FC III, Alper SL et al. A glucose transport protein expressed predominately in insulin-responsive tissues. *Proc Natl Acad Sci USA* 1989;86:2535-2539
255. Richter EA, Hargreaves M. Exercise, GLUT4, and skeletal muscle glucose uptake. *Physiol Rev* 2013;93:993-1017
256. Dagaard JR, Nielsen JN, Kristiansen S et al. Fiber type-specific expression of GLUT4 in human skeletal muscle: influence of exercise training. *Diabetes* 2000;49:1092-1095
257. Megeney LA, Neuffer PD, Dohm LG et al. Effects of muscle activity and fiber composition on glucose transport and GLUT4. *Am J Physiol Endocrinol Metab* 1993;264:E583-E593
258. Houmard JA, Shinebarger MH, Dolan PL et al. Exercise training increases GLUT-4 protein concentration in previously sedentary middle-aged men. *Am J Physiol Endocrinol Metab* 1993;264:E896-E901
259. Host HH, Hansen PA, Nolte LA et al. Rapid reversal of adaptive increase in muscle Glut-4 and glucose transport capacity after training cessation. *J Appl Physiol* 1998;84:798-802
260. Kranjic GN, Cameron-Smith D, Hargreaves M. Effect of short-term training on GLUT-4 mRNA and protein expression in human skeletal muscle. *Exp Physiol* 2004;89:559-563
261. Frosig C, Rose AJ, Trebak JT et al. Effects of endurance exercise training on insulin signaling in human skeletal muscle: interactions at the level of phosphatidylinositol 3-kinase, AKT, and AS160. *Diabetes* 2007;56:2093-2102
262. Greiwe JS, Hickner RC, Hansen PA et al. Effects of endurance exercise training on muscle glycogen accumulation in humans. *J Appl Physiol* 1999;87:222-226
263. Houmard JA, Egan PC, Neuffer PD et al. Elevated skeletal muscle glucose transporter levels in exercise-trained middle-aged men. *Am J Physiol* 1991;261:E437-E443
264. Frøsig CH, Rose AJ, Trebak JT et al. Effects of endurance exercise training on insulin signaling in human skeletal muscle. *Diabetes* 2007;56:2093-2103
265. Mikus CR, Oberlin J, Libla AM et al. Lowering physical activity impairs glycemic control in healthy volunteers. *Med Sci Sports Exerc* 2012;44(2):225-231
266. Babraj JA, Vollaard NBJ, Keast C et al. Extremely short duration high intensity interval training substantially improves insulin action in young healthy males. *BMC Endocrine Disorders* 2009;9:3, doi:10.1186/1472-6823-9-3
267. Lee D, Sui X, Church TS et al. Associations of cardiorespiratory fitness and obesity with risk of impaired fasting glucose and type 2 diabetes in men. *Diabetes Care* 2009;32:257-262
268. The Inter Act Consortium. Physical activity reduces the risk of incident type 2 diabetes in general and in abdominally lean and obese men and women: the EPIC-Inter Act Study. *Diabetologia* 2012;55:1944-1952
269. Andersen E, Ekelund U, Anderssen SA. Effects of reducing sedentary time on glucose metabolism in immigrant Pakistani men. *Med Sci Sports Exerc* 2015;47(4):775-781
270. Church TS, LaMonte MJ, Barlow CE et al. Cardiorespiratory fitness and body mass index as predictors of cardiovascular disease mortality among men with diabetes. *Arch Int Med* 2005;165(18):2114-2120
271. Zethelius B, Gudjónsdóttir S, Eliasson B et al. Level of physical activity is associated with risk of cardiovascular diseases and mortality in patients with type 2 diabetes: report from the Swedish National Diabetes Register. *Eur J Prevent Cardiol* 2014;21(2):244-251
272. Lee SS, Kang S. Effects of regular exercise on obesity and type 2 diabetes mellitus in Korean children: improvements of glycemic control and serum adipokines level. *J Phys Ther Sci* 2015;27:1903-1907
273. Demissie S, Levy D, Benjamin EJ et al. Insulin resistance, oxidative stress, hypertension, and leucocyte telomere length in men from the Framingham Heart Study. *Aging Cell* 2006;5:325-330

274. Adaikalakoteswari A, Balasubramanyam M, Mohan V. Telomere shortening occurs in Asian type 2 diabetic patients. *Diab Med* 2005;22(9):1151-1156
275. Su Y, Liu XM, Sun YM et al. The relationship between endothelial dysfunction and oxidative stress in diabetes and prediabetes. *Int J Clin Pract* 2008;62:877-872
276. Hansel B, Giral P, Nobecourt E et al. Metabolic syndrome is associated with elevated oxidative stress and dysfunctional dense high-density lipoprotein particles displaying impaired antioxidative activity. *J Clin Endocrinol Metab* 2004;89:4963-4971
277. Matthews C, Gorenne I, Scott S et al. Vascular smooth muscle cells undergo telomere based senescence in human atherosclerosis: effects of telomerase and oxidative stress. *Circ Res* 2006;99:156-164
278. Salpea KD, Talmud PJ, Cooper JA et al. Association of telomere length with type 2 diabetes, oxidative stress and UPC2 gene variation. *Atherosclerosis* 2010;209:42-50
279. Adaikalakoteswari A, Balasubramanyam M, Ravikumar R et al. Association of telomere shortening with impaired glucose tolerance and diabetic macroangiopathy. *Atherosclerosis* 2007;195:83-89
280. Verzola D, Gandolfo MT, Gaetani G et al. Accelerated senescence in the kidneys of patients with type 2 diabetic nephropathy. *Am J Physiol* 2008;295:F1563-F1573
281. Tentolouris N, Nzietchueng R, Catan V et al. White blood cells telomere length is shorter in males with type 2 diabetes and microalbuminuria. *Diabetes Care* 2007;30:2909-2915
282. Sampson MJ, Hughes DA. Chromosomal telomere attrition as a mechanism for the increased risk of epithelial and senescent phenotypes in type 2 diabetes. *Diabetologia* 2006;49:1726-1731
283. Olivieri F, Lorenzi M, Antonicelli R et al. Leucocyte telomere shortening in elderly type 2 DM patients with previous myocardial infarction. *Atherosclerosis* 2009;206:588-593
284. Uziel O, Singer JA, Danicek V et al. Telomere dynamics in arteries and mononuclear cells of diabetic patients: effect of diabetes and of glycemic control. *Exp Gerontol* 2007;42:971-978
285. Salpea KD, Humphries SE. Telomere length in atherosclerosis and diabetes. *Atherosclerosis* 2010;209:35-38
286. Laditka SD, Laditka JN. Active life expectancy of Americans with diabetes: risk of heart disease, obesity, and inactivity. *Diab Res Clin Pract* 2015;107:37-45
287. Kruk J, Aboul-Enen HY. Physical activity in the prevention of cancer. *Asian Pac J cancer Prev* 2006;7:11-21
288. Ma J, Pollak MN, Giovannucci E et al. Prospective study of colorectal cancer risk in men and plasma levels of insulin-like growth factor IGF-1 and IGF-binding protein 3. *J Natl Cancer Inst* 1999;91:620-625
289. Key TJ, Appleby PN, Reeves GK et al. Sex hormones and risk of breast cancer in premenopausal women: a collaborative reanalysis of individual participant data from seven prospective studies. *Lancet Oncol* 2013;14(10):1009-1019
290. Ilyasova D, Colbert LH, Harris TB et al. Circulating levels of inflammatory markers and cancer risk in the health aging and body composition cohort. *Cancer Epidemiol Biomarker Prev* 2005;14(10):2413-2428
291. Carmichael AR, Bates T. Obesity and breast cancer: a review of the literature. *The Breast* 2004;13:85-92
292. McTiernan AR. Intervention studies in exercise and cancer prevention. *Med Sci Sport Exerc* 2003;35:1841-1845
293. Hackney AC, Fahrnes CL, Gullledge TB. Basal reproductive hormonal profiles are altered in endurance trained men. *J Sports Med Phys Fitness* 1998;38:138-141
294. Leitzmann M, Powers H, Anderson AS et al. European Code against cancer 4th edition: physical activity and cancer. *Cancer Epidemiol* 2015; Jul 9, pii S 1877-7821(15)00127-7 doi:10.1016/j.canep.2015.05.009. Pub ahead of print

295. Shepard RJ, Shek PN. Association between physical activity and susceptibility to cancer. Possible mechanism. *Sport Med* 2002;26:293-315
296. Feig DI, Reid TM, Loeb LA. Reactive oxygen species in tumorigenesis. *Cancer Res* 1994;54:1890S-1894S
297. Chi F, Wu R, Zeng YC et al. Circulation insulin-like growth factor peptides and colorectal cancer risk: an updated systematic review and meta-analysis. *Mol Biol Rep* 2013;40(5):3583-3590
298. Li T, Wei S, Shi Y et al. The dose-response effect of physical activity on cancer mortality: findings from 71 prospective cohort studies. *Br J Sports Med* 2015, Sept 18. pii: bjsports-2015-094927. doi:10.1136/bj sports-2015-094927 (E pub ahead of print)
299. Shi Y, Li T, Wang Y et al. Household physical activity and cancer risk: a systematic review and dose-response meta-analysis of epidemiological studies. *Sci Rep* 5, 14901; doi 10.1038/s rep 14901 (2015)
300. Healy GN, Matthews CE, Dunstan DW et al. Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-6. *Eur Heart J* 2011;32(5):590-597
301. Shen D, Mao W, Liu T et al. Sedentary behaviour and incident cancer: a meta-analysis of prospective studies. *PLoS One* 2014;9(8).e105 709
302. Chao A, Connell JC, Jacobs EJ et al. Amount, type, and timing of recreational physical activity in relation to colon and rectal cancer in older adults: the Cancer Prevention Study II. Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* 2004;3(12):2187-2195
303. Je Y, Jeon JY, Giovannucci EL et al. Association between physical activity and mortality in colorectal cancer: a meta-analysis of prospective cohort studies. *Int J Cancer* 2013;133:1905-1913
304. Chen Y, Yu C, Li Y. Physical activity and risks of esophageal and gastric cancers: a meta-analysis. *PLoS One* 2014;9(2):e 88082
305. Vigen C, Bernstein L, Wu AH. Occupational physical activity and risk of adenocarcinomas of the esophagus and stomach. *Int J Cancer* 2006;118:1004-1009
306. Singh S, Varayil JE, Devanna S et al. Physical activity is associated with reduced risk of gastric cancer: a systematic review and meta-analysis. *Cancer Prev Res* 2015;7(1):12-22
307. Meyerhardt JA, Giovannucci EL, Holmes MD et al. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 2006;24(22):3527-3534
308. Meyerhardt JA, Giovannucci EL, Ogino S et al. Physical activity and survival in male colorectal cancer survivors. *Arch Intern Med* 2009;169(22):2102-2108
309. Meyerhardt JA, Heseltine D, Niedzwiecki D et al. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALBG 89803. *J Clin Oncol* 2006;24(22):3535-3541
310. Campbell PT, Patel AV, Newton CC et al. Association of recreational physical activity and leisure time spent sitting with colorectal cancer survival. *J Clin Oncol* 2013;31(7):876-885
311. Campbell PT, Newton CC, Dehal AN et al. Impact of body mass index on survival after colorectal cancer diagnosis: The Cancer Prevention Study II Nutrition Cohort. *J Clin Oncol* 2012;30(1):42-52
312. Christopoulos PF, Msaduel P, Koutsilies M. The role of insulin-like-growth factor-1 system in breast cancer. *Molecular Cancer* 2015;14:43- doi:10.1186/s12943-015-0291-7
313. Kaaks R, Johnson T, Tikk K et al. Insulin-like growth factor 1 and risk of breast cancer by age and hormone receptor status- A prospective study within the EPIC cohort. *Int J Cancer* 2014;134:2683-2690
314. Renehan AG, Tyson AG, Egger M et al. Body mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 2008; 371(9612):569-578
315. Patterson RE, Cadmus LA, Emond JA et al. Physical activity, diet, adiposity, and female breast cancer prognosis: a review of the epidemiologic literature. *Maturitas* 2010;66(1):5-15

316. Surmacz E. Leptin and adiponectin: emerging therapeutic targets in breast cancer. *J Mammary Gland Biol & Neoplasia* 2013;18:321-332
317. Coughlin SS, Smith SA. The insulin-like growth factor axis, adipokines, physical activity, and obesity in relation to breast cancer incidence and recurrence. *Cancer Clin Oncol* 2015; 14(2):24-31
318. Key T, Appleby P, Barnes I et al. Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 2002; 94(8):606-616
319. Yager JD, Davidson NE. Estrogen carcinogenesis in breast cancer. *N Engl J Med* 2006; 354(3):270-282
320. Carpenter CL, Duvall K, Jardack P et al. Weight loss reduces breast ductal fluid estrogens in obese postmenopausal women: a single arm intervention pilot study. *Nutr J* 2012;11:102, doi:10.1186/1475-2891-11-102
321. Jones ME, Schoemaker M, Rae M et al. Changes in estradiol and testosterone levels in postmenopausal women after changes in body mass index. *J Clin Endocrinol Metab* 2013; 98(7):2957-2974
322. Neilson HK, Conroy SM, Friedenreich CC. The influence of energetic factors on biomarkers of postmenopausal breast cancer risk. *Curr Nutr Rep* 2014;3:22-34
323. Van Gemert WA, Schuit AJ, van der Palen J et al. Effect of weight loss, with or without exercise, on body composition and sex hormones in postmenopausal women: the SHAPE-2 trial. *Breast Cancer Res* 2015;17:120, doi:10.1186/s13058-015-0633-9
324. Veljkovic M, Dopsaj V, Dopsaj M et al. Physical activity and natural anti-VIP antibodies: potential role in breast and prostate cancer therapy. *PLoS One* 2011;6(11):e28304
325. Moody TW, Leyton J, Gozes I et al. VIP and breast cancer. *Ann New York Acad Sciences* 1998;865:290-296
326. Moody TW, Gozes I. Vasoactive intestinal peptide receptors: a molecular target in breast and lung cancer, 2007; *Curr Pharmaceut Design* 2007;13:1099-1104
327. Valdehita A, Bajo AM, Schally AV et al. Vasoactive intestinal peptide (VIP) induces trans-activation of EGFR and HER2 in human breast cancer cells. *Molecular Cellular Endocrinol* 2009;302:41-48
328. Nagakawa O, Junicho OA, Akashi T et al. Vasoactive intestinal peptide and pituitary adenylate cyclase activating polypeptide stimulate interleukin-6 production in prostate cancer cells and prostatic epithelial cells. *Oncology Reports* 2005;13:1217-1221
329. Gkonos PJ, Ashby MH, Andrade AA. Vaso active intestinal peptide stimulates prostate-specific antigen secretion by LNCaP prostate cancer cells. *Regulatory Peptides* 1996;65:153-157
330. Paul S, Heinz-Erian P, Said SI. Autoantibody to vasoactive intestinal peptide in human circulation. *Biochem Biophys Res Commun* 1985;130:479-485
331. Delgado-Cruzata L, Wu HC, Perrin M et al. Global DNA-methylation levels in white blood cells-DNA from sisters discordant for breast cancer from the New York site of the Breast Cancer Family Registry. *Epigenetics* 2012;7(8):868-874
332. Ntanasis-stathopoulos J, Tzanninis JG, Philippou et al. Epigenetic regulation of gene expression induced by physical exercise. *J Musculoskelet Neuronal Interact* 2013;13(2):133-146
333. Zeng H, Irwin ML, Lu L et al. Physical activity and breast cancer survival: an epigenetic link through reduced methylation of a tumor suppressor gene L3MBTL1. *Breast Cancer Res Treat* 2012;133:127-135
334. White AJ, Sandler DP, Bolick SCE et al. Recreational and household physical activity at different time points and DNA global methylation. *Eur J Cancer* 2013;49(9):2199-2206
335. Body NF, Lockwood GA, Martn LJ et al. Mammographic densities and breast cancer risk. *Breast Dis* 1998;10:113-126

336. Tamimi RM, Byrne C, Colditz GA et al. Endogenous hormone levels, mammographic density, and subsequent risk of breast cancer in postmenopausal women. *J Natl Cancer Inst* 2007;99(15):1178-1187
337. Qureshi SA, Ellingjord-Dale M, Hofvind S et al. Physical activity and mammographic density in a cohort of postmenopausal Norwegian women; a cross-sectional study. *Springer PLUS* 2012;1:75, doi:10.1186/2193-2193-1801-1-75
338. Vachon CM, Kuni CC, Anderson K et al. Association of mammographically defined percent breast density with epidemiological risk factors for breast cancer (United States). *Cancer Causes Control* 2000;11:653-662
339. Yaghjian L, Colditz GA, Wolin K et al. Physical activity and mammographic breast density: a systemic review. *Breast Cancer Res Treat* 2012;135:367-380
340. Trinh T, Eriksson M, Darabi H et al. Background risk of breast cancer and the association between physical activity and mammographic density. *Breast Cancer Res* 2015;17:50
341. Chaddock-Heyman L, Mackenzie MJ, Zuniga K et al. Higher cardiorespiratory fitness levels are associated with greater hippocampal volume in breast cancer survivors. *Frontiers in Human Neuroscience* 2015;9:Article 465 doi:10.3389/fnhum.2015.00465
342. Argyriou AA, Assimakopoulos K, Iconomou G et al. Either called “chemobrain” or “chemofog” the longform chemotherapy-induced cognitive decline in cancer survivors is real. *J Pain Symptom Manage* 2011;41:126-139
343. Pereira-Dias G, Hollywood R, Bevilacqua MC et al. Consequences of cancer treatments on adult hippocampal neurogenesis: implications for cognitive function and depressive symptoms. *Neuro Oncol* 2014;16:476-492
344. De Rviter MB, Reneman L, Boogerd W et al. Late effects of high-dose adjuvant chemotherapy on white and gray matter in breast cancer survivors: converging results from multimodal magnetic resonance imaging. *Hum Brain Mapp* 2012;33:2971-2983
345. Dietrich J. Chemotherapy associated central nervous system damage. *Adv Exp Med Biol* 2010;678:77-85
346. Bergouignan L, Lefranc JP, Chupin M et al. Breast cancer affects both the hippocampus volume and the episodic autobiographical memory retrieval. *PLoS One* 2011;6:e25349
347. Kesler S, Janelins M, Koovakkattu D et al. Reduced hippocampal volume and verbal memory performance associated with interleukin-6 and tumor necrosis factor-alpha levels in chemo-therapy-treated breast cancer survivors. *Brain Behav Immun* 2013;30:S109-S116
348. Mustafa S, Werner M, Inagaki M et al. 5-Fluorouracil chemotherapy affects spatial working memory and newborn neurons in the adult rat hippocampus. *Eur J Neuroscience* 2008;28:323-330
349. Seigers R, Fardell JE. Neurobiological basis of chemotherapy-induced cognitive impairment: a review of rodent research. *Neurosci Biobehav Rev* 2011;35:729-741
350. Fardell JE, Vardy J, Shah JD et al. Cognitive impairments caused by oxaliplatin and 5-fluorouracil chemotherapy are ameliorated by physical activity. *Psychopharmacology (Berl)* 2012; 220:183-193
351. Naylor AS, Bull C, Nilsson MK et al. Voluntary running rescues adult hippocampal neurogenesis after irradiation of the young mouse brain. *Proc Acad Sci USA* 2008;105:14632-14637
352. Wong-Goodrich S, Pfau ML, Flores CT et al. Voluntary running prevents progressive memory decline and increases adult hippocampal neurogenesis and growth factor expression after whole-brain irradiation. *Cancer Res* 2010;70:9329-9338
353. Bugg JM, Head D. Exercise moderates age-related atrophy of the medial temporal lobe. *Neurobiol Aging* 2011;32:506-514
354. Chaddock L, Erickson KI, Prakash RS et al. A neuro imaging investigation of the association between aerobic fitness, hippocampal volume and memory performance in preadolescent children. *Brain Res* 2010;1358:172-183

355. Erickson KI, Prakash RS, Voss MW et al. Aerobic fitness is associated with hippocampal volume in elderly humans. *Hippocampus* 2009;19:1030-1039
356. Bernstein L, Patel AV, Ursin G et al. Lifetime recreational exercise activity and breast cancer risk among black women and white women. *J Natl Cancer Inst* 2005;97:1671-1679
357. Cohen SS, Matthews CE, Bradshaw PT et al. Sedentary behaviour, physical activity, and likelihood of breast cancer among black and white women: a report from the Southern Community Cohort Study. *Cancer Prev Res* 2013;6(6):566-576
358. Suzuki R, Iwasaki M, Yamamoto S et al. Leisure-time physical activity and breast cancer risk defined by estrogen and progesterone receptor status. The Japan Public Health Center-based Prospective Study. *Prev Med* 2011;52:227-233
359. Liu Y, Hu F, Li D et al. Does physical activity reduce the risk of prostate cancer? A systemic review and meta-analysis. *Eur Urol* 2011;60(5):1029-1044
360. Moore SC, Gierach GL, Schatzkin A et al. Physical activity, sedentary behaviours, and the prevention of endometrial cancer. *Br J Cancer* 2010;103(7):933-938
361. Leitzman MF, Koebnick C, Abnet CC et al. Prospective study of physical activity and lung cancer by histologic type in current, former, and never smokers. *Am J Epidemiol* 2009;169(5):542-553
362. Behrens G, Leitzman MF. The association between physical activity and renal cancer: systemic review and meta-analysis. *Brit J Cancer* 2013;108(4):798-81
363. Friedenreich CM, Orenstein MR. Physical activity and cancer prevention: etiologic evidence and biological mechanisms. *J Nutr* 2002;132:3456S-3464S
364. Sachdeva A, Kumar K, Anand KS. Non pharmacological cognitive enhancers-current perspectives. *J Clin Diagnost Res* 2015;9(7):VE01-VE06
365. APA (1994). American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, 4th Edn. Washington DC: American Psychiatric Assoc
366. Plassman BL, Langa KM, Fisher GG et al. Prevalence of dementia in the United States: the aging, demographics, and memory study. *Neuroepidemiology* 2007;29(1-2):125-132
367. Plassman BL, Langa KM, Fisher GG et al. Prevalence of cognitive impairment without dementia in the United States. *Ann Intern Med* 2008;148(6):427-434
368. ADI (2010) World Alzheimer Report. The Global Economic Impact of Dementia. London: Alzheimer's Disease International. Online: [http://www.alz.co.uk/research/files/World Alzheimer Report 2010 Executive Summary.pdf](http://www.alz.co.uk/research/files/World%20Alzheimer%20Report%202010%20Executive%20Summary.pdf)
369. Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's Disease prevalence. *Lancet Neurol* 2011;10:819-828
370. Etgen T, Sander D, Förstl H. Mild cognitive impairment and dementia: the importance of modifiable risk factors. *Deutsches Ärzteblatt International* 2011;108(44):743
371. Mattson MP, Maudsley S, Martin B. BDNF and 5HT: a dynamic duo in age-related memorial plasticity and neurodegenerative disorders. *Trends Neurosci* 2004;27:589-594
372. Karege F, Perret G, Bondolf G et al. Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Res* 2002;109:143-148
373. Krabbe KS, Nielsen AR, Kikkrogh-Madsen R et al. Brain derived neurotrophic factor (BDNF) and type 2 diabetes. *Diabetologica* 2007;50:431-438
374. Ebadi M, Bashir RM, Heidick ML et al. Neurotrophins and their receptors in nerve injury and repair. *Neurochem Int* 1997;30:347-374
375. Tyler WT, Alonso M, Bramham CR et al. From acquisition to consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-dependent learning. *Learn Mem* 2002;9:224-237
376. Cassilhas RC, Viana VA, Grassmann V et al. The impact of resistance exercise on cognitive function of the elderly. *Med Sci Sports Exerc* 2007;39(8):1401-1407

377. Trejo JL, Carro E, Torres-Alemán I. Circulating insulin-like growth factor 1 mediates exercise-induced increases in the number of new neurons in the adult hippocampus. *J Neurosci* 2001;21(5):1628-1634
378. Seifert T, Brassard P, Wissenberg M et al. Endurance training enhances BDNF release from the human brain. *Am J Physiol Regul Integr Comp Physiol* 2010;298:R372-R377
379. Adlard PA, Perreau VM, Pop V et al. Voluntary exercise decreases amyloid load in a transgenic model of Alzheimer's Disease. *J Neurosci* 2005;25(17):4217-4221
380. Van Praag H, Shubert T, Zhao C et al. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci* 2005;25(38):8680-8685
381. Penke L, Maniega SM, Murray C et al. A general factor of brain white matter integrity predicts information processing speed in healthy older people. *J Neurosci* 2010;30(22):7569-7574
382. Bherer L, Erickson KI, Liu-Ambrose T. A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J Aging Res* 2013; Article ID 657508 <http://dx.doi.org/10.1155/2013/657508>
383. Raz N, Ghiletta P, Rodrigue KM et al. Trajectories of brain aging in middle-aged and older adults: regional and individual differences. *Neuroimage* 2010;51(2):501-511
384. Fjell AM, Westlye LT, Grydeland H et al. Critical ages in the life course of adult brain: nonlinear subcortical aging. *Neurobiol Aging* 2013;34:2239-2247
385. Erickson KI, Raji CA, Lopez OL et al. Physical activity predicts gray matter volume in late adulthood. *Neurology* 2010;75:1415-1422
386. Colcombe SJ, Erickson KL, Scalf PE et al. Aerobic exercise training increases brain volume in aging humans. *J Gerontol* 2006;61A(11):1166-1170
387. Killgore WDS, Olson EA, Weber M. Physical exercise habits correlate with gray matter volume of the hippocampus in healthy adult humans. *Scientific Reports* 2013;3:3457 [doi:10.1038/srep03457](https://doi.org/10.1038/srep03457)
388. Erickson KI, Voss MW, Prakash RS et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci USA* 2011;108:3017-3022
389. Ten Brinke LF, Bolandzadeh N, Nagamatsu LS et al. Aerobic exercise increases hippocampal volume in older women with probable mild cognitive impairment: a 6-month randomized controlled trial. *Brit J Sports Med* 2015;49(4):248-254
390. Burns JM, Cronk BB, Anderson HS et al. Cardiorespiratory fitness and brain atrophy in early Alzheimer's Disease. *Neurology* 2008;71(3):210-216
391. Honea R, Thomas GP, Harsha A et al. Cardiorespiratory fitness and preserved medial temporal lobe volume in Alzheimer's Disease. *Alzheimer Dis Assoc Disord* 2009;23(3):188-197
392. Taaffe DR, Irie F, Masaki KH et al. Physical activity, physical function, and incident dementia in elderly men: The Honolulu-Asia Aging Study. *J Gerontol* 2008;63A(5):529-535
393. Larson EB, Wang L, Bowen JD et al. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann Int Med* 2006;144:73-81
394. Liu R, Sui X, Laditka JN et al. Cardiorespiratory fitness as a predictor of dementia mortality in men and women. *Med Sci Sports Exerc* 2012;44(2):253-259
395. Llamas-Velasco S, Contador I, Villarejo-Galende A et al. Physical activity as a protective factor against dementia: a prospective population-based study (NEDICES). *J Int Neurophysiol Soc* 2015;21(10):861-867
396. Sofi F, Valecchi D, Bacci D et al. Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J Inter Med* 2011;269(1):107-117
397. Middleton LE, Barnes DE, Liu LY et al. Physical activity over the life course and its association with cognitive performance and impairment in old age. *J Am Geriatr Soc* 2010;58(7):1322-1326

398. Liu-Ambrose T, Nagamatsu LS, Voss MW et al. Resistance training and functional plasticity of the aging brain: a 12-month randomized controlled trial. *Neurobiol Aging* 2012;33(8):1690-1698
399. Cassilas RC, Viana VA, Grassman V et al. The impact of resistance exercise on the cognitive function of the elderly. *Med Sci Sports Exerc* 2007;39(8):1401-1407
400. Baker LD, Frank LL, Foster-Schubert K et al. Effects of aerobic exercise on mild cognitive impairment: a controlled trial. *Arch Neurol* 2010;67(1):71-79
401. Kwak YS, Um S, Son TG et al. Effect of regular exercise on senile dementia patients. *Int J Sports Med* 2008;29(6):471-474
402. Teri L, Gibbons LE, McCurry SM et al. Exercise plus behavioral management in patients with Alzheimer Disease: a randomized controlled trial. *JAMA* 2003;290(15):2015-2022
403. Heyn P, Abreu BC, Ottenbacher KJ. The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil* 2004;85:1694-1704
404. Avadhani R, Fowler K, Barbato C et al. Glycemia and cognitive function in metabolic syndrome and coronary heart disease. *Am J Med* 2015;128:46-55
405. Jefferson AL. Cardiac output as a potential risk factor for abnormal brain aging. *J Alzheimer's Dis* 2010;20(3):813-821
406. Cermakova P, Eriksson M, Lund LH et al. Heart failure and Alzheimer's Disease. *J Internat Med* 2015;277:406-425
407. Gary RA, Brunn K. Aerobic exercise as an adjunct therapy for improving cognitive function in heart failure. *Cardiol Research and Practice* 2014; Article ID 157508
<http://dx.doi.org/10.1155/2014/157508>
408. Saczynski JS, Beiser A, Seshadri S et al. Depressive symptoms and risk of dementia. The Framingham Heart Study. *Neurology* 2010;75:35-41
409. Hickies I, Naismith S, Ward PB et al. Reduced hippocampal volumes and memory loss in patients with early- and late onset depression. *Brit J Psychiatry* 2005;186:197-202
410. Simon GE. Social and economic burden of mood disorders. *Biol Psychiatry* 2003;54:208-215
411. Blazer DG. Depression in late life: review and commentary. *J Gerontol A Biol Sci Med Sci* 2003;58:249-265
412. Rajkowska G. Cell pathology in mood disorders. *Semin Clin Neuropsychiatry* 2002;7:281-292
413. McEwen BS. Effects of adverse experiences for brain structure and function. *Biol Psychiatry* 2000;48:721-731
414. Ninan I. Synaptic regulation of affective behaviors; role of BDNF. *Neuropharmacology* 2014;76(00):doi:10.1016/j.neuropharm.2013.04.011
415. Erickson KI, Miller DL, Roecklein KA. The aging hippocampus: interactions between exercise, depression, and BDNF. *Neuroscientist* 2012;18(1):82-97
416. Dwivedi Y. Involvement of brain-derived neurotrophic factor in late-life depression. *Am J Geriatr Psychiatry* 2013;21(5):433-449
417. Claes S, Myint AM, Domschke K et al. The kynurenine pathway in major depression: haplotype analysis of three related functional candidate genes. *Psychiatry Res* 2011;188:355-360
418. Müller N, Schwarz MJ. The immune-mediated alteration of serotonin and glutamate: towards an integrated view of depression. *Mol Psychiatry* 2007;12:988-1000
419. Myint AM, Kim YK. Network beyond IDO in psychiatric disorders: revisiting neurodegeneration hypothesis. *Prog Neuropsychopharmacol Biol Psychiatry* 2014;48:304-313
420. Schwarcz R, Bruno JP, Muchowski PJ et al. Kynurenines in the mammalian brain: when physiology meets pathology. *Nat Rev Neurosci* 2012;13:465-477

421. Agudelo LZ, Femenia T, Orhan F et al. Skeletal muscle PGC-1 α 1 modulates kynurenine metabolism and mediates resilience to stress-induced depression. *Cell* 2014;159:33-45
422. Schmidt HD, Duman RS. Peripheral BDNF produces antidepressant-like effects in cellular and behavioural models. *Neuropsychopharmacology* 2010;35:2378-2391
423. Martinowich K, Lu B. Interaction between BDNF and serotonin: role in mood disorders. *Neuropsychopharmacology* 2008;33:73-83
424. Shirayama Y, Chen ACH, Nakagawa S et al. Brain-derived neurotrophic factor produces antidepressant effects in behavioural models of depression. *J Neurosci* 2002;22(8):3251-3261
425. Brunoni AR, Lopes M, Fregni F. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. *Int J Neuropsychopharmacol* 2008;11(8):1169-1180
426. Hassmén P, Koivula N, Uutela A. Physical exercise and psychological well-being: a population study in Finland. *Prevent Med* 2000;30:17-25
427. Sieverdes JC, Ray BM, Sui X et al. Association between leisure time physical activity and depressive symptoms in men. *Med Sci Sports Exerc* 2012;44(2):260-265
428. Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. *J Gerontol Med Sci* 1997;52A(1):M27-M35
429. Mather AS, Rodriguez C, Guthrie MF et al. Effects of exercise on depressive syndromes in older adults with poorly responsive depressive disorder. *Brit J Psychiatry* 2002;180:411-415
430. Babyak M, Blumenthal J, Herman S et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 month. *Psychosomat Med* 2000;62:633-638
431. Craft LL, Perna FM. Benefits of exercise for clinically depressed. *Prim Care Companion J Clin Psychiatry* 2004;6:104-111
432. United States Census Bureau. Survey of income and program participation. http://www.census.gov/people/disability/publications/disab_10/table_1.pdf
433. Federal interagency Forum on Aging Related Statistics. Older Americans key indicators of well-being. http://agingstats.gov/agingstats_net/Main_site/Data/2010_Documents/docs/Health_Status.pdf
434. Hennessy S, Kuruchi JE, Pan Q et al. Disability stage is an independent risk factor for mortality in medicare beneficiaries aged 65 years and older. *PM&R* 2015;7(12):1215-1225
435. Moreland JD, Richardson JA, Goldsmith CH et al. Muscle weakness and falls in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc* 2004;52:1121-1129
436. Ruiz JR, Sui X, Lobelo F et al. Association between muscular strength and mortality in men: a prospective cohort study. *BMJ* 2008;337:92-95
437. Frontera WR, Suh D, Krivickas LS et al. Skeletal muscle fiber quality in older men and women. *Am J Physiol Cell Physiol* 2000;279:C611-C618
438. De Brito LBB, Ricardo DR, De Araújo DSMS et al. Ability to sit and rise from the floor as a predictor of all-cause mortality. *Europ J Prev Cardiol* 2014;21(7):892-898
439. Kuh D. A life course approach to healthy aging, frailty, and capability. *J Gerontol A Biol Sci Med Sci* 2007;62A:717-721
440. Boyle PA, Buchman AS, Wilson RS et al. Physical frailty is associated with incident mild cognitive impairment in community-based older adults. *J Am Geriatr Soc* 2010;58(2):248-255
441. Petterson MJ, Giuliani C, Morey MC et al. Physical activity as a preventative factor for frailty: The Health, Aging, and Body Composition Study. *J Gerontol Med Sci* 2009;64A(1): 61-68
442. Rockwood K, Howlett SE, MacKnight C et al. Prevalence, attributes, and outcomes of fitness and frailty in community-dwelling older adults: report from the Canadian study of health and aging. *J Gerontol A Biol Sci Med Sci* 2004;59(12):1310-1317
443. El-Khoury F, Cassou B, Charles MA et al. The effects of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomized controlled trials. *BMJ* 2013;347:f6234

444. Helbostad JL, Sletvold O, Moe-Nilssen R. Home training with and without additional group training in physically frail old people living at home: effect on health-related quality of life and ambulation. *Clin Rehabil* 2004;18:498-508
445. Aguirre LE, Villareal DT. Physical exercise as therapy for frailty. *Nestle Nutr Workshop Ser* 2015;83:83-92
446. De Martinis M, Franceschi C, Monti D et al. Inflammation markers predicting frailty and mortality in the elderly. *Exp Mol Pathol* 2006;80:219-227
447. Brüünsgaard H, Pedersen BK. Age-related inflammatory cytokines and disease. *Immunol Allergy Clin North Am* 2003;23(1):15-39
448. Zhang Q, Raouf M, Chen Y et al. Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* 2010;464:104-107
449. Oka T, Hikoso S, Yamaguchi O et al. Mitochondrial DNA that escapes from autophagy causes inflammation and heart failure. *Nature* 2012;485:251-255
450. Pinti M, Cevenini E, Nasi M et al. Circulating mitochondrial DNA increases with age and is a familiar trait: implications for “inflamm-aging”. *Brit J Immunol* 2014;44:1552-1562
451. Nasi M, Cristani A, Pinti M et al. Decreased circulating mtDNA levels in professional male volleyball players. *Int J Sports Physiol Perform* 2015 Jun 10 (Epub ahead of print)
452. Traustadóttir T, Davies SS, Su Y et al. Oxidative stress in older adults: effects of physical fitness. *AGE* 2012;34:969-982
453. Ashok BT, Ali R. The aging paradox: free radical theory of aging. *Exp Gerontol* 1999;34:293-303
454. Butterfield DA, Howard B, Yatin S et al. Elevated oxidative stress in models of normal brain aging and Alzheimer’s disease. *Life Sci* 1999;65(18-19):1883-1892
455. Spector A. Review: oxidative stress and disease. *J Oncol Pharmacol* 2000;16:193-201
456. Radak Z, Chung HY, Hisashi N et al. Age-associated increases in oxidative stress and nuclear transcription factor κ B activation are attenuated in rat liver by regular exercise. *FASEB J* 2004;18(6):749-750
457. Rosa EF, Silva AC, Ihara SS et al. Habitual exercise programme protects murine intestinal, skeletal, and cardiac muscles against aging. *J Appl Physiol* 2005;99(4):1569-1575
458. Cogley JN, Sakellariou GK, Murray S et al. Lifelong endurance training attenuates age-related genotoxic stress in human skeletal muscle. *Longevity & Healthspan* 2013;2:11-18
459. Blair SN, Kohl HW, Barlow CE et al. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 1995;273:1093-1108
460. Castillo-Garzon MJ, Ruiz JR, Ortega FB et al. Anti-aging therapy through fitness enhancement. *Clin Intern Aging* 2006;1:213-220
461. Rantanen T, Masaki K, He Q et al. Midlife muscle strength and human longevity up to age 100 years: a 44-year prospective study among a decedent cohort. *Age* 2012;34:563-570
462. Semba RD, Cappola AR, Sun K et al. Plasma klotho and mortality risk in older community-dwelling adults. *J Gerontol A Biol Sci Med Sci* 2011;66:794-800
463. Xiao NM, Zhang YM, Zheng Q et al. Klotho is a serum factor related to human aging. *Chin Med J* 2004;117:742-747
464. Semba RD, Cappola AR, Sun K et al. Relationship of low plasma klotho with poor grip strength in older community-dwelling adults: the InCHIANTI Study. *Europ J Appl Physiol* 2012; 112:1215-1220
465. Crasto CL, Semba RD, Sun K et al. Relationship of low-circulating “anti-aging” klotho hormone with disability in activities of daily living among older community-dwelling adults. *Rejuvenation Res* 2012;15:295-301
466. Kuro O M, Matsumura Y, Aizawa H et al. Mutation of the mouse klotho gene leads to a syndrome resembling aging. *Nature* 1997;390:45-51
467. Avin KG, Coen PM, Huang W et al. Skeletal muscle as a regulator of the longevity protein, Klotho. *Frontiers in Physiol* 2014;5:Article 189;doi:10.3389/fphys.2014.00189

468. Kurosu H, Yamamoto M, Clark JD et al. Suppression of aging in mice by the hormone Klotho. *Science* 2005;309:1829-1833
469. Brack S, Conboy IM, Roy S et al. Increased Wnt signaling during aging alters muscle stem cell fate and increases fibrosis. *Science* 2007;317:807-810
470. Doi S, Zou Y, Togao O et al. Klotho inhibits transforming growth factor β 1 (TGF- β 1) signaling and suppresses renal fibrosis and cancer metastasis in mice. *J Biol Chem* 2011;286:8655-8665
471. Lin J, Wu H, Tarr PT et al. Transcriptional co-activator PGC-1 α drives the formation of slow-twitch muscle fibres. *Nature* 2002;418(6899):797-801
472. Lai CH, Ho TJ, Kuo WW et al. Exercise training enhanced SIRT 1 longevity signaling replaces the IGF 1 survival pathway to attenuate aging-induced rat heart apoptosis. *AGE* 2014;36:9706 doi 10.1007/s11357-014-9706-4
473. Kwak HB, Song W, Lawler JM. Exercise training attenuates age-induced elevation in BAX/Bcl-2 Ratio, apoptosis, and remodeling in the rat heart. *FASEB J* 2006;20:791-793
474. Zhang JH, Zhang Y, Herman B. Caspases, apoptosis and aging. *Ageing Res Rev* 2003;2:357-366
475. Soufi FC, Farajnia S, Aslanabadi N et al. Long-term exercise training effects age-induced changes in HSP70 and apoptosis in rat heart. *Gen Physiol Biophys* 2008;27:263-270
476. Kwak HB. Effects of aging and exercise training on apoptosis in the heart. *J Exercise Rehabil* 2013;9(2):212-219
477. Levine HJ. Rest heart rate and life expectancy. *JACC* 1997;30(4):1104-1106
478. Coburn AF, Grey RM, Rivera SM. Observations on the relation of heart rate, life span, weight and mineralization in the digoxin-treated A/J mouse. *Johns Hopkins Med J* 1971;128:169-193
479. Stressman J, Jacobs JM, Stressman-Lande I et al. Aging, resting pulse rate, and longevity. *J Am Geriatr Soc* 2013;61(1):40-45
480. Zhang D, Shen X, Qi X. Resting heart rate and all-cause and cardiovascular mortality in the general population: a meta-analysis. *CMAJ* 2015; doi:10.1503/cmaj.150535 Publ. ahead of print
481. Nanchen D, Leening JGM, Locatelli I et al. Resting heart rate and the risk of heart failure in healthy adults: The Rotterdam Study. *Circ Heart Fail* 2013;6:403-410
482. Della Valle E, Stranges S, Trevisan M, et al. Self-rated measures of physical activity and cardiovascular risk in a sample of Southern Italian male workers: the Olivetti heart study. *Nutr Metab Cardiovasc Dis* 2004;14:143-149
483. Black A, Murray LJ, Cardwell C et al. Secular trends in heart rate in young adults, 1949 to 2004: analysis of cross-sectional studies. *Heart* 2006;92:468-473
484. De Souza CA, Shapiro LP, Clevenger CM et al. Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 2000;102:1351-1357
485. Marijon E, Tafflet M, Antero-Jaquemin J et al. Mortality of french participants in the Tour de France (1947-2012). *Eur Heart J* 2013;34:3145-3150
486. Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality: systematic review and dose-response meta-analysis of cohort studies. *Int J Epidemiol* 2011;40:1382-1400
487. Lee D, Pate RR, Lavie CJ et al. Leisure-time running reduces all-cause and cardiovascular mortality risk. *J Am Coll Cardiol* 2014;64(5):472-481
488. Kokkinos P, Myers J, Faselis C et al. Exercise capacity and mortality in older men. A 20-year follow-up study. *Circulation* 2010;122:790-797
489. Hamer M, Lavoie KL, Bacon SL. Taking up physical activity in later life and healthy aging: the English longitudinal study of aging. *Br J Sports Med* 2014;48:239-243

490. Zhao W, Ukawa S, Kawamura T et al. Health benefits of daily walking on mortality among younger-elderly men with or without major critical diseases in the New Integrated Suburban Seniority Investigation Project: a prospective cohort study. *J Epidemiol* 2015;25(19):609-616
491. Rizzuto D, Orsini N, Qiu C et al. Lifestyle, social factors and survival after age 75: population-based study. *BMJ* 2012;345:e5568 doi:10.1136/bmj.e5568
492. Wen CP, Wai JP, Tsai MK et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011; 378(9798): 1244-1253
493. Farrell SW, Finley CE, Haskell WL et al. Is there a gradient of mortality risk among men with low cardiorespiratory fitness? *Med Sci Sports Exerc* 2015;47(9):1825-1832
494. Arem H, Moore SC, Patel A et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. *JAMA Intern Med* 2015;175(6):959-967
495. Ruiz JR, Sui X, Lobelo F et al. Association between muscular strength and mortality in men: prospective cohort study. *BMJ* 2008;337:92-95; 2008;337:a439
496. Booth FW, Laye MJ, Roberts MD. Lifetime sedentary living accelerates some aspects of secondary aging. *J Appl Physiol* 2011;111:1497-1504
497. Prévost A, Lafitte M, Pucheu Y et al. Education and home-based training for intermittent claudication: functional effects and quality of life. *Eur J Prevent Cardiol* 2015;22(3):373-379
498. Kang JJE, Bodary PF. How old are your arteries? Exercise-mediated protection from age-associated vascular stiffness. *J Am Heart Assoc* 2014;3:e000941 doi:10.1161/JAHA.114.000941
499. Mitchell GF. Arterial stiffness and hypertension. *Hypertension* 2014;64:13-18
500. Jefferis BJ, Whincup PH, Papacosta O et al. Protective effect of time spent walking on risk of stroke in older men. *Stroke* 2014;45:194-199
501. Ridgel AL, Walter BL, Tatsuoka C et al. Enhanced exercise therapy in Parkinson's disease: a comparative effectiveness trial. *J Science Med Sport* 2016;19:12-17
502. Kirsten DK, Taube C, Lehnigk B et al. Exercise training improves recovery in patients with COPD after an acute exacerbation. *Respiratory Medicine* 1998;92:1191-1198
503. Kokkinos P, Faselis C, Myers JM et al. Exercise capacity and risk of chronic kidney disease in US veterans: a cohort study. *Mayo Clin Proc* 2015;90(4):461-468
504. Sawant A, House AA, Overend TJ. Anabolic effect of exercise training in people with end-stage renal disease on hemodialysis: a systematic review with meta-analysis. *Physiotherapy Canada* 2014;66(1):44-53
505. Position paper on transplantation and physical activity. T0066, PA/PH/TO[14]33:2 July 2015
506. Clarke SF, Murphy EF, O'Sullivan O et al. Exercise and associated dietary extremes impact on gut microbial diversity. *GUT* (Jun, 2014), doi:10.1136/gutjul-2013-306541
507. Berman S, Petriz B, Kajèniene A et al. The microbiota: an exercise immunology perspective. *Exerc Immunol Rev* 2015;21:70-79
508. Haskell WL, Lee IM, Pate RR et al. Physical activity and public health. Updated recommendation for adults from the American College of Sports Medicine and The American Heart Association. *Circulation* 2007;116:1081-1093
509. Eckel RH, Jakicic JM, Ard JD et al. 2013 AH/ACC Guideline on lifestyle management to reduce cardiovascular risk. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;129(Suppl 2):S76-S99
510. Gamber CE, Blissmer B, Deschenes MR et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43(7):1334-1359
511. Warburton DER, Nicol CW, Bredin SSD. Prescribing exercise as preventive therapy. *CMAJ* 2006;174(7):961-973

512. The American Heart Association Recommendations for Physical Activity in Kids: www.heart.org/Kids Activity Recommendations
513. Am College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 2009;41(3):687-708
514. Chodzko-Zajko WJ, Proctor DN, Fiatarone S et al. American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc* 2009;41(7):1510-1530
515. Fletcher GF, Ades PA, Kligfield P et al. Exercise standards for testing and training. A scientific statement from the American Heart Association. *Circulation* 2013;128:873-934
516. Borjesson M, Umhauen A, Kouidi E et al. Cardiovascular evaluation of middle-aged/ senior individuals engaged in leisure-time sport activities: position stand from the sections of exercise physiology and sports cardiology of the European Association of cardiovascular prevention and rehabilitation. *Eur J Cardiovasc Prev Rehab* 2011;8(3):446-458
517. Pearson SJ, Young A, Macaluso A et al. Muscle function in elite master weight lifters. *Med Sci Sports Exerc* 2002;34(7):1199-1206
518. Tanaka H, Seales DR. Invited review: Dynamic exercise performance in masters athletes: insight into the effects of primary human aging on physiological functional capacity. *J Appl Physiol* 2003;95:2152-2162
519. Piepoli MF, Corras U, Adamopoulos A et al. Secondary prevention in the clinical management of patients with cardiovascular diseases. Core components, standards and outcome measures for referral and delivery. *Eur J Prev Cardiol* 2014;21(6):664-681
520. Mezzani A, Hamm LF, Jones AM et al. Aerobic exercise intensity assessment and prescription in cardiac rehabilitation: a joint position statement of the European Association for Cardiovascular Prevention and Rehabilitation, the American Association of Cardiovascular and Pulmonary Rehabilitation and the Canadian Association of Cardiac Rehabilitation. *Eur J Prev Cardiol* 2012;20(3):442-467
521. Kulik A, Ruel M, Jneid H et al. Secondary prevention after coronary artery bypass graft surgery: a scientific statement from the American Heart Association. *Circulation* 2015;131:927-964
522. Wise FM. Exercise-based cardiac rehabilitation in chronic heart failure. *Austral Fam Physician* 2007;3(12):1019-1024
523. Dubach P, Sixt S, Myers J. Exercise training in chronic heart failure: why, when, and how. *Swiss Med Wkly* 2001;131:510-514
524. Ades PA. Cardiac rehabilitation exercise and self care for chronic heart failure. *JACC Heart Fail* 2013;1(6):540-547
525. De Maeyer C, Beckers P, Vrints CJ et al. Exercise training in chronic heart failure. *Ther Adv Chronic Dis* 2013;4(3):105-117
526. Working Group on Cardiac Rehabilitation & Exercise Physiology and Working Group on Heart Failure of the European Society of Cardiology. Recommendations for exercise training in chronic heart failure patients. *Eur Heart J* 2001;22:125-135
527. Lampert R, Olshansky B, Heidbuchel H et al. Safety of sports for athletes with implantable cardioverter-defibrillators. Results of a prospective, multinational registry. *Circulation* 2013;127:2021-2030
528. Kramer DB, Mitchell SL, Monteiro J et al. Patient activity and survival following implantable cardioverter-defibrillator implantation: The ALTITUDE Activity Study. *J Am Heart Assoc* 2015;4:e001775
529. Heidbuchel H, Carre F. Exercise and competitive sports in patients with an implantable cardioverter-defibrillator. *Eur Heart J* 2014;35:3097-3102
530. Corrado D, Schmied C, Basso C et al. Risk of sports: do we need a pre-participation screening for competitive and leisure athletes? *Eur Heart J* 2011;32:934-944

531. Maron BJ, Thompson PD, Ackerman MJ et al. Recommendations and considerations related to preparticipation screening for cardiovascular abnormalities in competitive athletes: 2007 Update. *Circulation* 2007;115:1643-1655
532. Pelliccia A, Fagard R, Bjørnstad HH et al. Recommendations for competitive sports participation in athletes with cardiovascular disease. A consensus document from the Study Group of Sports Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Disease of the European Society of Cardiology. *Eur Heart J* 2005;26:1422-1445
533. Corrado D, Pelliccia A, Bjørnstad HH et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. *Eur Heart J* 2005;26:516-524
534. Thompson PD, Franklin BA, Balady GJ et al. Exercise and acute cardiovascular events. Placing the risks into perspective. A scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism and the Council on Clinical Cardiology. *Circulation* 2007;115:2358-2368
535. Stecker EC, Reinier K, Marijon E et al. Public health burden of sudden cardiac death in the United States. *Circ Arrhythm Electrophysiol* 2014;7:212-217
536. Marijon E, Tafflet M, Celermajer DS et al. Sports-related sudden death in the general population. *Circulation* 2011;124:672-681
537. Marijon E, Uy-Evanado A, Reinier K et al. Sudden cardiac arrest during sports activity in middle age. *Circulation* 2015;131:1384-1391
538. Harmon KG, Drezner JA, Maleszewski JJ et al. Pathogenesis of sudden cardiac death in National Collegiate Athletic Association. *Circ Arrhythm Electrophysiol* 2014;7:198-204
539. Harmon KG, Asif IM, Maleszewski JJ et al. Incidence, cause, and comparative frequency of sudden cardiac death in national collegiate athletic association. *Circ* 2015;132:10-19
540. Department of Health and Aging. Get up and Grow: Healthy Eating and Physical Activity for Early Childhood. Physical activity guidelines for 0-5 years old. 2009. [http://www.health.gov.au/internet/main/publishing.nsf/Content/EDFEDB588460BCE3CA25762B00232A13/\\$File/gug-directorscoord-2.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/EDFEDB588460BCE3CA25762B00232A13/$File/gug-directorscoord-2.pdf)
541. American Alliance for Health PE, Recreation and Dance (AAH-PERD). NASPE releases first ever physical activity guidelines for infants and toddlers. *Illinois J Health PE Rec Dance* 2002;50:31-42
542. Hinkley T, Salmon J et al. Preschoolers' physical activity, screen time and compliance with recommendations. *Med Sci Sports Exerc* 2012;44(3):458-465
543. Trost SG, Mc Coy TA, Vanderveur SS et al. Physical activity patterns of inner-city elementary schoolchildren. *Med Sci Sports Exerc* 2013;45(3):470-474
544. Center for Disease Control and Prevention. Youth risk behaviour surveillance-United States. 2011. *MMWR* 2012;61(4)
545. Townsend N, Bhatnagar P, Wickramasinghe K et al. Physical activity statistics 2012, London, British Heart Foundation 2012
546. Martin K, Rosenberg M, Miller M et al. Move and munch final report. In: Trends in physical activity, nutrition, and body size in Western Australian children and adolescents: the Child and Adolescent Physical Activity and Nutrition Survey (CAPANS) 2008
547. Metcalf BS, Hosking J, Jeffery AN. Exploring the adolescent fall in physical activity: A 10-yr Cohort Study. *Med & Science Sports & Exercise* 2015;47(10):2084-2092
548. Wall MI, Carlson SA, Stein AD et al. Trends by age in youth physical activity: Youth Media Campaign Longitudinal Study. *Med Sci Sports Exerc* 2011;43(11):2140-2147
549. Ng SW, Popkin BM. Time use and physical activity: a shift away from movement across the globe. *Obesity reviews* 2012;13:659-680
550. Jefferis BJ, Sartini C, Ash S et al. Trajectories of objectively measured physical activity in free-living older men. *Med Sci Sports Exerc* 2015;47(2):343-349

551. Keevil VL, Wijndaele K, Luben R et al. Television viewing, walking speed, and grip strength in a prospective cohort study. *Med Sci Sports Exerc* 2015;47(4):735-742
552. Dür W, Griebler R (2007). Die Gesundheit der österreichischen SchülerInnen im Lebenszusammenhang. Ergebnisse des WHO-HBSC-Survey 2006. Wien: Bundesministerium für Gesundheit
553. Fonds Gesundes Österreich. Österreichische Empfehlungen für gesundheitswirksame Bewegung Band Nr. 8 aus der Reihe WISSEN
554. European Commission. Eurobarometer 72.3-Sport and Physical Activity; 2010 // in *Cancer Epidemiology* 2015;395:S46-S55. Leitzmann M, Powers H, Anderson AS et al. European code against cancer 4th edition: physical activity and cancer
555. Matthews CE, Moore SC, Sampson J et al. Mortality benefits for replacing sitting time with different physical activities. *Med Sci Sports Exerc* 2015;47(9):1833-1840
556. Lee IM, Shiroma J, Lobelo F et al. Effect of physical inactivity on major non-communicable diseases world wide: an analysis of burden of disease and life expectancy. *Lancet* 2012;380(9838):219-229
557. Flegal KM, Carroll MD, Ogden CL et al. Prevalence and trends in obesity among US adults 1999-2000. *JAMA* 2002;288:1723-1727
558. Cutler JA, Sorlie PD, Wolz M et al. Trends in hypertension prevalence, awareness, treatment, and control rates in United States adults between 1988-1994 and 1999-2004. *Hypertension* 2008;52:818-827
559. Wilmot KA, O'Flaherty M, Capewell S et al. Coronary heart disease mortality declines in the United States from 1979 through 2011. Evidence for stagnation in young adults, especially women. *Circulation* 2015;132:997-1002
560. Geiss LS, Wang J, Cheng YJ et al. Prevalence and incidence trends for diagnosed diabetes among adults aged 20 to 79 years, United States 1980-2012. *JAMA* 2014;312:1218-1226
561. Hulsege G, Picavet HSJ, Blokstra A et al. Today's adult generation are less healthy than their predecessors: generation shifts in metabolic risk factors: the Doetinchem Cohort Study. *Eur J Prev Cardiol* 2014;21(9):1134-1144
562. Pratt M, Macera CA, Wang G. Higher direct medical costs associated with physical inactivity. *The Physician and Sportsmedicine* 2000;28(10):63-70
563. World Health Organization. Global Action Plan for the Prevention and Control of Noncommunicable Diseases: 2013-2020. Geneva, Switzerland: WHO Press: 2013, <http://www.who.int/global-coordination-mechanism/publications/global-action-plan-ncds-eng.pdf>
564. Atun J, Jaffar S, Nishtar S et al. Improving responsiveness of health systems to non-communicable diseases. *Lancet* 2013;381(9867):690-697
565. Centers for Disease Control and Prevention, Chronic Disease Prevention and Health Promotion. <http://www.cdc.gov/chronic-disease/overview/index.htm>
566. Ng M, Fleming T, Robinson M et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384(9945):766-778
567. Mc Kinsey Global Institute. *Overcoming Obesity: an initial economic analysis*. New York, NY: Mc Kinsey and Company 2014
568. Arena R, Guazzi M, Lianov L et al. Healthy lifestyle interventions to combat noncommunicable disease—a novel nonhierarchical connectivity model for key stakeholders: a policy statement from the American Heart Association, European Society of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and American College of Preventive Medicine. *Mayo Clin Proc* 2015;90(8):1082-1103
569. Dumith SC, Hallal PC, Reis RS et al. Worldwide prevalence of physical inactivity and its association with human development index in 76 countries. *Prev Med* 2011;53(1-2):24-28
570. Kohl III HW, Craig CL, Lambert EV et al. The pandemic of physical inactivity: global action of public health. *Lancet* 2012;380(9838):294-305

571. Arena R, Lavie CJ. The healthy lifestyle team is central to the success of accountable care organizations. *Mayo Clin Proc* 2015;90(5):572-576
572. Bloom DE, Cafiero ET, Jané-Llopis E et al. The global economic burden of non-communicable diseases. Geneva: World Economic Forum; 2011
<http://www.hsph.harvard.edu/program-on-the-global-demography-of-aging/working-papers/2012/PGDA-WP-87.pdf>
573. Mozaffarian D, Benjamin G, Go AS et al. American Heart Association Statistical Committee and Stroke Statistics Subcommittee: heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation* 2015;131:e29-e322
574. Fox KM, Wang L, Gandra SR et al. Clinical and economic burden associated with cardiovascular events among patients with hyperlipidemia: a retrospective cohort study. *BMC Cardiovasc Disorders* 2016;13:1-15, doi:10.1186/s12872-016-0190-x
575. Ben-Yehuda O, Kazi DS, Bonafede M et al. Angina and associated healthcare costs following percutaneous coronary intervention: a real-world analysis from a multi-payer database. *Catheter Cardiac Interv* 2016, Jan 17 (E pub ahead of print); doi:10.1002/ccd.26365
576. Caruba T, Chevreul K, Zarca K et al. Annual costs of stable coronary artery disease in France: a modeling study. *Arch Cardiovasc Dis* 2015;108(11):576-588
577. Asaria M, Walker S, Palmer S et al. Using electronic health records to predict costs and outcomes in stable coronary artery disease. *Heart* 2016;0:1-8 (E pub ahead of print) doi:10.1136/heartj-2015-308850
578. Häußler J. Effects of obesity and physical activity on health care utilization and costs. University of Konstanz, Department of Economics. Working Paper Series 2014-07; <http://www.uni-konstanz.de/econdoc/working-paper-series/>
579. American Diabetes Association. Economic costs of diabetes in the U.S. in 2012. *Diabetes Care* 2013; 36:1033-1046
580. HEALTH-Institut für Biomedizin und Gesundheitswissenschaften der JOANNEUM RESEARCH Forschungsgesellschaft GmbH, im Auftrag von Sanofi-Aventis GmbH. Diabetes Forum, Sonderbeilage zur Ausgabe 1/2015
581. Wimo A, Winblad B, Jönsson L. The worldwide societal costs of dementia: estimates for 2009. *Alzheimer Dement* 2010;6:98-103
582. Wimo A, Jönsson L, Gustavsson A et al. The economic impact of dementia in Europe in 2008—cost estimates from the Eurocode project. *Int J Geriatric Psychiatry* 2011;26:825-832
583. Wübker A, Zwakhalen SM, Challis D et al. Costs of care for people with dementia just before and after nursing home placement: primary data from eight European countries. *Eur J Heart Econ* 2015;16(7):689-707
584. Farre M, Hard JM, Kostov B et al. Direct and indirect costs and resource use in dementia care: a cross-sectional study in patients living at home. *Int J Nurs Stud* 2016;55:39-49
585. Kelley AS, McGarry K, Gorges R et al. The burden of health care costs for patients with dementia in the last 5 years of life. *Ann Intern Med* 2015;163(10):729-736
586. Ziegler U, Doblhammer G. Prevalence and incidence of dementia in Germany—a study based on data from the public sick funds in 2002. *Gesundheitswesen* 2009;71(5):281-290
587. Scharzkopf LM, Menn P, Leidl R et al. Excess costs of dementia disorders and the role of age and gender—an analysis of German health and long-term care insurance claims data. *BMC Health Services Research* 2012;12:165
588. Roux L, Pratt M, Tengs TO et al. Cost effectiveness of community-based physical activity interventions. *Am J Prev Med* 2008;35(6):578-588
589. Allender S, Foster C, Scarborough P et al. The burden of physical activity-related ill health in the UK. *J Epidemiol Community Health* 2007;61:344-348
590. Stadt Wien (Hrsg). Bereichsleitung für Gesundheitsplanung und Finanzmanagement 2001, Gesundheit in Wien, Wiener Gesundheits- und Sozialsurvey Wien

591. World Health Organization. WHO 2006, Gesundheitsförderung durch Bewegung-ein Handlungsrahmen für die Europäische Region der WHO-Schritte zu einem körperlich aktiverem Europa. WHO-Regionalbüro für Europa Kopenhagen
592. Allender S, Rayner M. The burden of overweight and obesity-related ill health in the UK. *Obesity Reviews* 2007;8:467-473
593. Ghayoumi A, Myers J, Ablla J et al. Physical activity patterns and health care costs [abstract]. *J Am Coll Cardiol* 2004;45:16A
594. Bertram O. Men's Health. Das Workout ohne Geräte. Südwest Verlag ISBN 978-3-517-08922-5
595. Steinbach M. Women's Health. Das Workout ohne Geräte. Südwest Verlag ISBN 978-3-517-08992-8
596. Lauren M, Clark J. Fit ohne Geräte. Riva Verlag ISBN 978-3-86883-166-5
597. Blaschke TF, Osterberg I, Vrijens et al. Adherence to medications: insight arising from studies on the unreliable link between prescribed and actual drug dosing histories. *Ann Rev Pharmacol Toxicol* 2012;52:275-301
598. Sallis RES. Exercise is medicine and physicians need to prescribe it! *Br J Sports Med* 2009; 43(1):3-4
599. Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol* 2012;2(2):1143-1211
600. Löllgen H, Halle M, Wilhelm M et al. Sport auf Rezept-trotz hoher Evidenz vernachlässigt die Therapie körperliche Aktivität. *Cardio News Austria* 5.2015/3. Jahrgang, Abdruck Deutsches Ärzteblatt
601. Hung R, Al-Mallah AL, Qadi MA et al. Cardiorespiratory fitness attenuates risk for major adverse cardiac events in hyperlipidemic men and women independent of statin therapy: The Henry Ford Exercise Testing Project. *Am Heart J* 2015;170:390-399
602. Naci H, Ioannidis JPA. Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *Br J Sports Med* 2015;49:1414-1422
603. Hippocrates . The Aphorisms of Hippocrates; Birmingham, AL: The Classics of Medicine Library, Division of Gryphon Editions Ltd 1982
604. Aigelsreiter H. Die 7 Aigelsreiter, Eigenverlag, 5. Auflage 2006, ISBN 3-9501544-0-X
605. Frankl VE, Handl C. Bergerlebnis und Sinnerfahrung, Tyrolia-Verlag, 2. Auflage 2003, ISBN 3-7022-2466-1
606. Deslandes A. The biological clock keeps ticking, but exercise may turn it back. *Arq Neuropsiquiatr* 2013;72(2):113-118
607. Carroll DD, Courtney-Long EA, Stevens AC et al. Vital signs: disability and physical activity—United States, 2009-2012. *MMWR Morb Mortal Wkly Rep* 2014;63(18):407-413
608. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults *The Lancet* 2017;390, No 10113, p 2627-2642
609. Kivimäki M, Kuosma E, Ferrie JE et al. Overweight, obesity, and risk of cardiometabolic multimorbidity: pooled analysis of individual-level data for 120 813 adults from 16 cohort studies from the USA and Europe. *The Lancet Public Health* 2017;2, No 6, e 277-e 285, June
610. Holzweber F. Der volkswirtschaftliche Nutzen des Sports. Eine Dokumentation 2017